arXiv:1303.4863v1 [hep-ph] 20 Mar 2013

KIAS-P13015

Leptons and Quarks from a Discrete Flavor Symmetry

Y. H. Ahn*

School of Physics, KIAS, Seoul 130-722, Korea

(Dated: March 21, 2013)

Abstract

We propose a new model of leptons and quarks based on the discrete flavor symmetry T', the double covering of A_4, in which the hierarchies of charged fermion masses and the mildness of neutrino masses are responsible for Higgs scalars. After spontaneous breaking of flavor symmetry, with the constraint of renormalizability in the Lagrangian, the leptons have $m_e = 0$ and the quarks have the Cabibbo-Kobayashi-Maskawa (CKM) mixing angles $\theta^q_{12} = 13^\circ, \theta^q_{23} = 0^\circ$ and $\theta^q_{13} = 0^\circ$. Thus, certain effective dimension-5 operators are introduced, which induce $m_e \neq 0$ and lead the quark mixing matrix to the CKM one in form. On the other hand, the neutrino Lagrangian still keeps renormalizability. For completeness, we show numerical analysis: in the lepton sector, only normal mass hierarchy is permitted within 3σ experimental bounds with the prediction of both large deviations from maximality in the atmospheric mixing angle θ_{23} and the measured values of reactor angle. So, future precise measurements of θ_{23}, whether $\theta_{23} \to 45^\circ$ or $|\theta_{23} - 45^\circ| \to 5^\circ$, will either exclude or favor our model. Together with it, our model makes predictions for the Dirac CP phase, which is almost compatible with the global analysis in 1σ experimental bounds. Moreover, we show the effective mass $|m_{ee}|$ measurable in neutrinoless double beta decay to be in the range $0.04 \lesssim |m_{ee}|[eV] < 0.11$, which can be tested in near future neutrino experiments.

PACS numbers:

---

* Email: yhahn@kias.re.kr

I. INTRODUCTION

In the Standard Model (SM) of particle physics with a single Higgs there are enormously various hierarchies of quark and lepton Yukawa couplings, *that is*, $m_t/m_u = y_t/y_u \sim 10^5, m_\tau/m_e = y_\tau/y_e \sim 10^3$ etc.. In addition, there are several fundamental questions in Yukawa sector: why the top quark is uniquely so big compared with the other fermions, why neutrino masses are so small and so mild compared with the other charged fermions, why both the three leptonic mixing angles and one quark mixing angle are large, while the two quark mixing angles are so small. In some sense, our understanding of fermion masses and mixing angles remains at a very primitive level. One of the approaches often adopted in the understanding for a possible solution for the flavor puzzle consists of the introduction of family symmetries which constrain the flavor structure of Yukawa couplings and lead to predictions for fermion masses and mixings. We propose a simplified way to address those questions in a non-Abelian discrete symmetry T' [1], by both introducing six types of Higgs fields and imposing all Yukawa couplings being of order one. Then the group T' can be responsible for the present Pontecorvo-Maki-Nakagawa-Sakata (PMNS) and Cabibbo-Kobayashi-Maskawa (CKM) mixing matrices. And the hierarchies of fermions can originate from the different Higges. The representations of T' are those of A_4 plus three independent doublets $\mathbf{2}, \mathbf{2}', \mathbf{2}''$. Similar to A_4 [2–6] [1], in neutrino sector using four in-equivalent representations $\mathbf{1}, \mathbf{1}', \mathbf{1}''$ and $\mathbf{3}$ one can obtain the tri-bimaximal (TBM) [7] mixing pattern. In the presence of the doublet representations one can naturally describe the mass hierarchy among the charged fermions and the Cabibbo angle in the CKM matrix [8].

After the relatively large reactor angle θ_{13} measured in Daya Bay and RENO [9] including Double Chooz, T2K and MINOS experiments [10], the most recent analysis based on global fits [11] of the neutrino oscillations enters into a new phase of precise determination of mixing angles and mass squared differences, indicating that the TBM mixing for three flavors should be corrected in the lepton sector: their allowed ranges at 1σ (3σ) from global fits are given by

$$\theta_{13} = 8.66^{\circ}{}^{+0.44^{\circ}\ (+1.30^{\circ})}_{-0.46^{\circ}\ (-1.47^{\circ})}\ , \qquad \delta_{\rm CP} = 300^{\circ}{}^{+66^{\circ}\ \ (+60^{\circ})}_{-138^{\circ}\ (-300^{\circ})}\ , \qquad \theta_{12} = 33.36^{\circ}{}^{+0.81^{\circ}\ (+2.53^{\circ})}_{-0.78^{\circ}\ (-1.27^{\circ})}\ ,$$
$$\theta_{23} = 40.0^{\circ}{}^{+2.1^{\circ}}_{-1.5^{\circ}} \oplus 50.4^{\circ}{}^{+1.3^{\circ}}_{-1.3^{\circ}}\ 1\sigma, \quad (35.8^{\circ} \sim 54.8^{\circ}\ 3\sigma)\ ,$$

---

[1] The finite group A_4 describes the even permutations of four objects and possesses.

$$\Delta m^2_{\rm Sol}[10^{-5}{\rm eV}^2] = 7.50^{+0.18\ (+0.59)}_{-0.19\ (-0.50)}\ ,\ \Delta m^2_{\rm Atm}[10^{-3}{\rm eV}^2] = \begin{cases} 2.473^{+0.070\ (+0.222)}_{-0.067\ (-0.197)}\ , & {\rm NMH} \\ 2.427^{+0.042\ (+0.185)}_{-0.065\ (-0.222)}\ , & {\rm IMH} \end{cases} \quad (1)$$

where $\Delta m^2_{\rm Sol} \equiv m_2^2 - m_1^2$, $\Delta m^2_{\rm Atm} \equiv m_3^2 - m_1^2$ for the normal mass hierarchy (NMH), and $\Delta m^2_{\rm Atm} \equiv |m_3^2 - m_2^2|$ for the inverted one (IMH). While there are the large values of the solar mixing angle $\theta_{\rm sol} \equiv \theta_{12}$, the atmospheric mixing angle $\theta_{\rm atm} \equiv \theta_{23}$ and the reactor mixing angle $\theta_{\rm reac} \equiv \theta_{13}$ in the lepton sector, in the quark sector the Cabibbo angle and the other two small quark mixing angles, e.g., at 1σ level [12] read:

$$\theta^q_{12} = (13.03 \pm 0.05)^\circ\ , \quad \theta^q_{23} = (2.37^{+0.05}_{-0.09})^\circ\ , \quad \theta^q_{13} = (0.20^{+0.02}_{-0.02})^\circ\ , \quad \delta^q_{CP} = (67.17^{+2.78}_{-2.44})^\circ\ . (2)$$

The discrepancy of the mixing angles in Eqs. (1) and (2) may tell us about some new flavor symmetries of quarks and leptons. And it is well known that the mass spectrum of the charged fermions exhibits a strong hierarchical pattern [see Eq. (22)], unlike that of neutrinos which shows a mild hierarchy. These facts may provide a clue to the nature of quark-lepton physics beyond the SM [2]. Therefore, it is very important to find a natural model that leads to the observed flavor mixing patterns for quarks and leptons. In the present article we shall build such a model to emphasize the leptonic mixing parameters and the quarks one.

In this work we propose a new model based on flavor symmetry T' that can accommodate quarks and leptons in the same framework invariant under $SU(2)_L \times U(1)_Y \times T'$. We introduce six types of Higgses in the Yukawa sector to depict the mass hierarchies of fermions, and simultaneously the mixing parameters of the leptons and those of the quarks. In addition, in order to simplify our model and to remove the unwanted Yukawa terms appearing in the Lagrangian, we impose a continuous global symmetry which cannot be gauged. After spontaneous $U(1)_X$ breaking, to avoid Goldstone bosons it has to be explicitly broken down to a subgroup. We stress that in our model CP invariance is originally explicitly broken in the Lagrangian level by the complex Clebsch-Gordan coefficients, even though all parameters in Lagrangian being real are imposed. The lepton sector in our model can not only naturally explain large deviations from the TBM but also provide a possibility for low-energy CP violation in neutrino oscillations and the mildness of neutrino masses.

---

[2] The dada (1) and (2) at 1σ seem to disfavor the maximal mixing in the atmospheric mixing angle, indicating that it starts to disfavor the Quark-Lepton Complementarity [13] on $\theta^q_{23} + \theta_{23} = 45^\circ$, even it is not significant yet.

The paper is organized as follows. In the next section, we present the particle content together with the flavor symmetry of our model and the mass terms of both neutrino and charged fermion sectors after flavor symmetry breaking. In Sec. III, we show the neutrino masses generated in a type seesaw-I and their mixing angles and CP violation as well as the CKM matrix. In Sec. IV, for completeness, we show numerical analysis in the lepton sector. Then, we give the conclusion in Sec. IV, and briefly mention about the VEV alignments and spontaneous CP violation, as an example, in Appendix.

II. THE MODEL

In the absence of flavor symmetries, particle masses and mixings are generally undetermined in a gauge theory. Here, we present a discrete symmetry model based on a T' flavor symmetry for leptons and quarks in order to depict the mass hierarchies of charged fermions, the mildness of neutrino masses, and simultaneously the present mixing parameters of the neutrino oscillation data and those of the quarks. Moreover, we describe the model to understand CP violations in the lepton sector which is imperative, if the baryon asymmetry of the Universe (BAU) originated from leptogenesis scenario in the seesaw models [14].

Here we recall that T' is the symmetry group of the double tetrahedron [1, 15]. The group T' has 24 elements and has two kinds of representations. It contains the representations of A_4: one triplet $\mathbf{3}$ and three singlets $\mathbf{1}, \mathbf{1}'$ and $\mathbf{1}''$. When working with these representations there is no distinction between the group T' and the group A_4. In particular, in these representations, the elements of T' coincide in pairs and can be described by the same matrices that represent the elements in A_4. The other representations are three boublets $\mathbf{2}, \mathbf{2}'$ and $\mathbf{2}''$. The representations $\mathbf{1}', \mathbf{1}''$ and $\mathbf{2}', \mathbf{2}''$ are complex conjugated to each other. Note that A_4 is not a subgroup of T', since the two-dimensional representations can not be decomposed into representations of A_4. The generators S and T satisfy the relation $S^2 = R, T^3 = (ST)^3 = R^2 = \mathbf{1}$, where $R = \mathbf{1}$ in case of the odd-dimensional representation and $R = -\mathbf{1}$ for $\mathbf{2}, \mathbf{2}', \mathbf{2}''$ such that R commutes with all elements of the group. In the three-dimensional unitary representation, there are abelian subgroups of T' : Z_3, Z_4 and Z_6

symmetries, which are generated by the elements

$$T=\begin{pmatrix}1&0&0\\0&\omega&0\\0&0&\omega^2\end{pmatrix},\qquad TST^2=\frac{1}{3}\begin{pmatrix}-1&2&2\\2&-1&2\\2&2&-1\end{pmatrix},\qquad S=\frac{1}{3}\begin{pmatrix}-1&2\omega&2\omega^2\\2\omega^2&-1&2\omega\\2\omega&-1&2\omega^2\end{pmatrix},\tag{3}$$

respectively, where $\omega=e^{i2\pi/3}$ is a complex cubic-root of unity. Especially, the elements T and TST^2 are of importance for the structure of our model. The group T' has seven irreducible representations, one triplet $\mathbf{3}$, three doublets $\mathbf{2},\mathbf{2}',\mathbf{2}''$ and three singlets $\mathbf{1},\mathbf{1}',\mathbf{1}''$ with the multiplication rules $\mathbf{1}^j\otimes\mathbf{r}^k=\mathbf{r}^k\otimes\mathbf{1}^j=\mathbf{r}^{j+k}$ for $\mathbf{r}=1,2$, $\mathbf{1}^j\otimes\mathbf{3}=\mathbf{3}\otimes\mathbf{1}^j=\mathbf{3}$, $\mathbf{2}^j\otimes\mathbf{2}^k=\mathbf{3}\oplus\mathbf{1}^{j+k}=\mathbf{r}^{j+k}$, $\mathbf{2}^j\otimes\mathbf{3}=\mathbf{3}\otimes\mathbf{2}^j=\mathbf{2}\oplus\mathbf{2}'\oplus\mathbf{2}''$ and $\mathbf{3}\otimes\mathbf{3}=\mathbf{3}_s\oplus\mathbf{3}_a\oplus\mathbf{1}\oplus\mathbf{1}'\oplus\mathbf{1}''$, where $j,k=0,\pm1$ and we have denoted $\mathbf{1}^0\equiv\mathbf{1},\mathbf{1}^1\equiv\mathbf{1}',\mathbf{1}^{-1}\equiv\mathbf{1}''$ and similarly for the doublet representations. The sum $j+k$ is modulo 3. The Clebsch-Gordan coefficients for the decomposition of the product representations are shown in Ref. [15, 16].

We extend the standard model (SM) by the inclusion of an T'-triplet of right-handed $SU(2)_L$-singlet Majorana neutrinos N_R, and the introduction of six types of scalar Higgs fields: the SM $SU(2)_L$-doublet Higgs bosons Φ,Ψ, which we take to be T'-triplets $\mathbf{3}$ representation, the other two $SU(2)_L$-doublet Higgs bosons H,G, which are distinguished from Φ,Ψ by being T'-doublets $\mathbf{2}$ representation, and another $SU(2)_L$-doublet $\mathbf{2}$ of Higgs boson η, which is a T'-singlet $\mathbf{1}$ representation, finally an $SU(2)_L$-singlet T'-triplet $\mathbf{3}$ Higgs field χ:

$$\begin{aligned}\Phi_j&=\begin{pmatrix}\varphi_j^+\\\varphi_j^0\end{pmatrix},&\qquad H_k&=\begin{pmatrix}\phi_k^+\\\phi_k^0\end{pmatrix},&\qquad &\chi_j,\\ \Psi_j&=\begin{pmatrix}\psi_j^+\\\psi_j^0\end{pmatrix},&\qquad G_k&=\begin{pmatrix}G_k^+\\G_k^0\end{pmatrix},&\qquad \eta&=\begin{pmatrix}\eta^+\\\eta^0\end{pmatrix}\end{aligned}\tag{4}$$

where $j=1,2,3$ and $k=1,2$. According to the above Higgs scalars, we impose T' flavor symmetry for leptons and quarks. And, we levy all Yukawa couplings are of order one which implies that all the hierarchies of fermions appearing in the Lagrangian are responsible for the Higgses we introduced economically. In addition, after spontaneous breaking of flavor symmetry, the VEVs (vacuum expectation values) of such fields need spontaneous CP violation. The representations of the field content of the model under $SU(2)\times U(1)\times T'$ are summarized in Table-I, where each flavor of lepton doublets is assigned to one of the three T'-singlet representations: the electron-flavor to the $\mathbf{1}$, the muon flavor to the $\mathbf{1}''$, and the tau flavor to the $\mathbf{1}'$, and Q_L denotes left handed quark $SU(2)_L$ doublet and $\tau_R, \mathcal{E}_R$ and $t_R, \mathcal{U}_R$ $(b_R, \mathcal{D}_R)$ are the respective SM right handed lepton and u-type (d-type) $SU(2)_L$ singlets, respectively. Here, the down-type fermions $\mathcal{E}_R$, $\mathcal{D}_R$ and the up-type fermions $\mathcal{U}_R$ are assigned as T'-doublets and right handed gauge singlets:

$$\mathcal{E}_R \left\{ \begin{array}{l} e_R \\ \mu_R \end{array} \right. , \qquad \mathcal{D}_R \left\{ \begin{array}{l} d_R \\ s_R \end{array} \right. , \qquad \mathcal{U}_R \left\{ \begin{array}{l} u_R \\ c_R \end{array} \right. . \tag{5}$$

In the presence of three T'-triplet Higgs scalars Φ, Ψ, χ and two T'-doublets Higgs scalars H, G, Higgs potential Lagrangian involving interaction terms among Φ, χ, H, G, Ψ and χ, which would be written as $V(\Phi\chi)$, $V(\Phi H), V(\Phi G), V(\Psi H), V(\Psi G)$, $V(H\chi), V(G\chi)$ and $V(\Psi\chi)$, would be problematic for vacuum stability. Such stability problems can be naturally solved, for instance, in the presence of extra dimensions or in supersymmetric dynamical completions [6]. In these cases, those interaction terms are either disallowed or highly suppressed. In our model, the T' flavor symmetry is spontaneously broken by those T'-triplet and doublet scalars and T'-singlet scalar. From the condition of the global minima of the scalar potential, we can obtain vacuum alignments of the fields χ, Φ, Ψ, H, G relevant to achieve our goal. The Higgs potential of our model contains many terms, which is listed in Appendix A, Eqs. (A3)-(A9). We spontaneously break the T' flavor symmetry by giving non-zero vacuum expectation values to some components of the T'-triplets χ, Ψ and Φ. We take the T' symmetry breaking scale to be above the electroweak scale in our model, *i.e.*, $\langle\chi\rangle > \langle\Phi^0\rangle$. As seen in Appendix A, the minimization of our scalar potential gives the following vacuum expectation values (VEVs):

$$\begin{aligned} \langle\chi\rangle &= v_\chi e^{i\varphi}(1,1,1) , \qquad \langle\Phi^0\rangle = \frac{v_\Phi e^{i\gamma}}{\sqrt{2}}(1,0,0) , \qquad \langle\Psi^0\rangle = \frac{v_\Psi e^{i\zeta}}{\sqrt{2}}(1,0,0) , \\ \langle\eta^0\rangle &= \frac{v_\eta}{\sqrt{2}} , \qquad \langle H^0\rangle = \frac{1}{\sqrt{2}}(v_{H_1}e^{i\rho_1}, v_{H_2}e^{i\rho_2}) , \qquad \langle G^0\rangle = \frac{1}{\sqrt{2}}(v_{G_1}e^{i\sigma_1}, v_{G_2}e^{i\sigma_2}) . \end{aligned} \tag{6}$$

The SM VEV $v = (\sqrt{2}G_F)^{-1/2} = 246$ GeV results from the combination $v^2 = \Sigma_k(v_\eta^2 + v_\Psi^2 + v_{H_k}^2 + v_{G_k}^2 + v_\Phi^2)$ where $k = 1, 2$. The non-zero expectation value $\langle\chi\rangle \sim (1,1,1)$ breaks T' symmetry down to a residual Z_4 symmetry which is generated by the group element TST^2. The non-zero expectation values $\langle\Phi^0\rangle \sim (1,0,0)$ and $\langle\Psi^0\rangle \sim (1,0,0)$ break T' symmetry down to its subgroup Z_3 which is generated by the group element T. The non-zero expectation value $\langle\eta^0\rangle = v_\eta/\sqrt{2}$ does not break the T' symmetry, because it is T'-flavorless. The non-zero expectation values $\langle H^0\rangle$ and $\langle G^0\rangle$ break $T' \rightarrow$ *nothing* with

TABLE I: Representations of the fields under T' and $SU(2)_L \times U(1)_Y$.

Field	L_τ, L_μ, L_e	Q_L	$\tau_R, \mathcal{E}_R$	$t_R, \mathcal{U}_R$	$b_R, \mathcal{D}_R$	N_R	χ	η	Φ	H	Ψ	G
T'	$\mathbf{1}'$, $\mathbf{1}''$, $\mathbf{1}$	$\mathbf{3}$	$\mathbf{1}'$, $\mathbf{2}''$	$\mathbf{1}'$, $\mathbf{2}'$	$\mathbf{1}'$, $\mathbf{2}''$	$\mathbf{3}$	$\mathbf{3}$	$\mathbf{1}$	$\mathbf{3}$	$\mathbf{2}$	$\mathbf{3}$	$\mathbf{2}$
$SU(2)_L \times U(1)_Y$	$(2, -\frac{1}{2})$	$(2, \frac{1}{6})$	$(1, -1)$	$(2, \frac{2}{3})$	$(2, -\frac{1}{3})$	$(1, 0)$	$(1, 0)$	$(2, \frac{1}{2})$	$(2, \frac{1}{2})$	$(2, \frac{1}{2})$	$(2, \frac{1}{2})$	$(2, \frac{1}{2})$

hierarchical breakings. In addition to T' flavor symmetry, we impose an additional symmetry $U(1)_X$ which is continuous global symmetry, where L_e, L_μ, L_τ, τ_R, $\mathcal{E}_R$, b_R and $\mathcal{D}_R$ carry $X = 1$ and Φ, G carry $X = -1$, while all other fields have $X = 0$. So this non-flavor symmetry forbids some irrelevant the $SU(2)_L \times U(1)_Y \times T'$ invariant Yukawa terms from the Lagrangian (see later). And Since Goldstone bosons resulting from spontaneous $U(1)_X$ breaking via $\langle\Phi\rangle, \langle G\rangle \neq 0$ are not allowed phenomenologically, so the additional symmetry $U(1)_X$ has to be explicitly broken [3] down to a subgroup Z_2 under which $\Phi \to -\Phi, G \to -G$ and the fields L_e, L_μ, L_τ, τ_R, $\mathcal{E}_R$, b_R and $\mathcal{D}_R$ also switch sign.

In our Lagrangian, we assume that there is a cutoff scale Λ, above which there exists unknown physics.

A. The neutrino sector

The Yukawa interactions ($d \leq 5$) in the neutrino sector invariant under $SU(2)_L \times U(1)_Y \times T'$ can be written as

$$\begin{aligned} -\mathcal{L}^{\nu}_{\rm Yuk} &= y^{\nu}_1 \bar{L}_\tau (\tilde{\Phi} N_R)_{\mathbf{1}'} + y^{\nu}_2 \bar{L}_\mu (\tilde{\Phi} N_R)_{\mathbf{1}''} + y^{\nu}_3 \bar{L}_e (\tilde{\Phi} N_R)_{\mathbf{1}} \\ &+ \frac{1}{2} M (\overline{N^c_R} N_R)_{\mathbf{1}} + \frac{1}{2} y^{\nu}_R (\overline{N^c_R} N_R)_{\mathbf{3}_s} \chi + \text{h.c.}, \end{aligned} \tag{7}$$

where $\tilde{\Phi} \equiv i\tau_2 \Phi^*$ and τ_2 is a Pauli matrix. Note here that there are no dimension-5 operators driven by χ field in the neutrino sector, and the above Lagrangian in neutrino sector is renormalizable. In this Lagrangian, each flavor of neutrinos has its own independent Yukawa term, since they belong to different singlet representations $\mathbf{1}'$, $\mathbf{1}''$, and $\mathbf{1}$ of T': the neutrino Yukawa terms involve the T'-triplets Φ and N_R, which combine into the appropriate singlet representation. The right-handed neutrinos have an additional Yukawa term that involves the T'-triplet SM-singlet Higgs χ. The mass term $\frac{1}{2}M(\overline{N_R^c}N_R)_{\mathbf{1}}$ for the right-handed neutrinos is necessary to implement the seesaw mechanism by making the right-handed neutrino mass parameter M large. The additional symmetry $U(1)_R$, as shown before, guarantees that the $SU(2)_L \times U(1)_Y \times T'$ invariant Yukawa terms $\bar{L}_{e,\mu,\tau}\tilde{\Psi}N_R$ are forbidden from the Lagrangian.

---

[3] In Appendix A, there are interaction terms $(\Phi^\dagger\Psi)(\Phi^\dagger\Psi), (G^\dagger H)(G^\dagger H)$ which break explicitly $U(1)_X$ to remove the unwanted Goldstone bosons in the low energy spectrum.

After the breaking of the flavor and electroweak symmetries, with the VEV alignments as in Eq. (6) the Dirac neutrino and right-handed neutrino mass terms from the Lagrangian (7) result in

$$\begin{aligned}
-\mathcal{L}_m^\nu &= \frac{v_\Phi e^{i\gamma}}{\sqrt{2}}\left(y_3^\nu \bar{\nu}_e N_{R1} + y_2^\nu \bar{\nu}_\mu N_{R3} + y_1^\nu \bar{\nu}_\tau N_{R2}\right) \\
&+ \frac{M}{2}(\overline{N_{R1}^c}N_{R1} + \overline{N_{R2}^c}N_{R3} + \overline{N_{R3}^c}N_{R2}) + \frac{y_R^\nu v_\chi e^{i\varphi}}{6}\Big\{2\overline{N_{R1}^c}N_{R1} + 2\overline{N_{R2}^c}N_{R2} + 2\overline{N_{R3}^c}N_{R3} \\
&- \overline{N_{R2}^c}N_{R3} - \overline{N_{R3}^c}N_{R2} - \overline{N_{R1}^c}N_{R2} - \overline{N_{R2}^c}N_{R1} - \overline{N_{R1}^c}N_{R3} - \overline{N_{R3}^c}N_{R1}\Big\} + \text{h.c.}\ . \qquad (8)
\end{aligned}$$

Then, the neutrino Dirac mass terms and the right-handed Majorana neutrino mass terms are expressed as

$$m_D = \frac{v_\Phi e^{i\gamma}}{\sqrt{2}}\begin{pmatrix} y_3^\nu & 0 & 0 \\ 0 & 0 & y_2^\nu \\ 0 & y_1^\nu & 0 \end{pmatrix} \qquad (9)$$

$$M_R = \begin{pmatrix} M + \frac{2}{3}y_R^\nu v_\chi e^{i\varphi} & -\frac{1}{3}y_R^\nu v_\chi e^{i\varphi} & -\frac{1}{3}y_R^\nu v_\chi e^{i\varphi} \\ -\frac{1}{3}y_R^\nu v_\chi e^{i\varphi} & \frac{2}{3}y_R^\nu v_\chi e^{i\varphi} & M - \frac{1}{3}y_R^\nu v_\chi e^{i\varphi} \\ -\frac{1}{3}y_R^\nu v_\chi e^{i\varphi} & M - \frac{1}{3}y_R^\nu v_\chi e^{i\varphi} & \frac{2}{3}y_R^\nu v_\chi e^{i\varphi} \end{pmatrix}, \qquad (10)$$

where v_Φ, $y_{1,2,3}^\nu$ and M, y_R^ν, v_χ are real positive variables.

B. Charged fermion sector

In the charged fermion sector, the Yukawa interactions ($d \le 5$) including dimension-5 operators driven by the χ field, invariant under $SU(2)_L \times U(1)_Y \times T'$, are given by

$$\mathcal{L}_{\text{Yuk}}^f = \mathcal{L}_{\text{Yuk}}^d + \mathcal{L}_{\text{Yuk}}^u + \mathcal{L}_{\text{Yuk}}^\ell\ , \qquad (11)$$

where

$$\begin{aligned} -\mathcal{L}^{d}_{\rm Yuk} &= y_b(\bar{Q}_L\Psi)_{\mathbf{1}''}b_R + Y_d\bar{Q}_L(H\mathcal{D}_R)_{\mathbf{3}} \\ &+ \frac{y_b^{a(s)}}{\Lambda}[(\bar{Q}_L\Psi)_{\mathbf{3}}\chi]_{\mathbf{1}''}b_R + \frac{Y_1^d}{\Lambda}\bar{Q}_L(H\mathcal{D}_R)_{\mathbf{1}''}\chi + \frac{Y_d^{a(s)}}{\Lambda}\bar{Q}_L(H\mathcal{D}_R)_{\mathbf{3}}\chi + \text{h.c.}, \qquad (12) \\ -\mathcal{L}^{u}_{\rm Yuk} &= y_t(\bar{Q}_L\tilde{\Phi})_{\mathbf{1}''}t_R + Y_u\bar{Q}_L(\tilde{G}\mathcal{U}_R)_{\mathbf{3}} \\ &+ \frac{y_t^{a(s)}}{\Lambda}[(\bar{Q}_L\tilde{\Phi})_{\mathbf{3}}\chi]_{\mathbf{1}''}t_R + \frac{Y_1^u}{\Lambda}\bar{Q}_L(\tilde{G}\mathcal{U}_R)_{\mathbf{1}'}\chi + \frac{Y_u^{a(s)}}{\Lambda}\bar{Q}_L(\tilde{G}\mathcal{U}_R)_{\mathbf{3}}\chi + \text{h.c.}\ , \qquad (13) \\ -\mathcal{L}^{\ell}_{\rm Yuk} &= y_\tau\bar{L}_\tau\eta\ \tau_R + Y_\mu\bar{L}_\mu(H\mathcal{E}_R)_{\mathbf{1}''} \\ &+ \frac{Y_1^\mu}{\Lambda}\bar{L}_\mu[(H\mathcal{E}_R)_{\mathbf{3}}\chi]_{\mathbf{1}''} + \frac{Y_2^\mu}{\Lambda}\bar{L}_e[(H\mathcal{E}_R)_{\mathbf{3}}\chi]_{\mathbf{1}} + \text{h.c.}, \qquad (14) \end{aligned}$$

with $\tilde{G} \equiv i\tau_2 G^*$. In the charged-lepton sector tau lepton involves the T'-singlet η and the T'-singlet right-handed charged-lepton τ_R, while muon lepton has the T'-doublet H and the T'-doublet right-handed charged-lepton $\mathcal{E}_R$ and there is no corresponding electron lepton term in renormalizable terms, indicating directly electron mass can be generated by the dimension-5 operators driven by χ field. And the τ-mass is generated upon the breaking of $T'-$invariant. Thus, the third family in charged leptons is different from the two, muon and electron. On the other hand, in the quark sectors the use of T'-doublets H, G and triplets Φ, Ψ Higgses can allow the third family to differ from the first two, and thus make plausible the mass hierarchies $m_t \gg m_c \gg m_u$ and $m_b \gg m_s \gg m_d$ [8]. And in the renormalizable terms the b-quark and t-quark masses are generated upon the breaking of $T' \to Z_3$. Mass terms of the quarks have two independent Yukawa terms with different couplings (y_b, Y_d) and (y_t, Y_u) for down-type quark and up-type quark, respectively, all involving T'-triplet Higgs fields Φ, Ψ and doublet fields H, G. The T'-triplet Φ is shared by both the three neutrino Yukawa terms and top-quark Yukawa term. The T'-doublet H is involved by the terms associated to the T'-doublet right-handed lepton $\mathcal{E}_R$ and down-type quark $\mathcal{D}_R$. As mentioned before, the above Yukawa Lagrangian has the additional symmetry $U(1)_X$. This non-flavor symmetry (continuous Global symmetry) ensures that the $SU(2) \times U(1) \times T'$ invariant Yukawa terms $\bar{Q}_L(G\mathcal{D}_R)_{\mathbf{3}}$,$\bar{Q}_L(G\mathcal{E}_R)_{\mathbf{3}}$, $\bar{Q}_L(\tilde{H}\mathcal{U}_R)_{\mathbf{3}}$, $(\bar{Q}_L\Phi)_{\mathbf{1}''}b_R$ and $(\bar{Q}_L\tilde{\Psi})_{\mathbf{1}''}t_R$ are absent from the Lagrangian.

In the charged fermion sectors from the Lagrangian (11), after the breaking of the flavor and electroweak symmetries, with the VEV alignments as in Eq. (6) the up-type quark and down-type quark mass terms result in

$$\mathcal{L}^{f}_{m} = \mathcal{L}^{d}_{m} + \mathcal{L}^{u}_{m} + \mathcal{L}^{\ell}_{m}\ , \qquad (15)$$

where

$$\begin{aligned}
-\mathcal{L}^d_m &= y_b\frac{\tilde{v}_\Psi}{\sqrt{2}}\bar{b}_L b_R + \frac{Y_d}{\sqrt{2}}\Big\{ i\tilde{v}_{H_1}\bar{d}_L d_R + \frac{1-i}{2}\bar{s}_L\left(d_R\tilde{v}_{H_2} + s_R\tilde{v}_{H_1}\right) + \tilde{v}_{H_2}\bar{b}_L s_R\Big\} \\
&+ \frac{\tilde{v}_\chi}{\sqrt{2}\Lambda}\Big\{ m^d_{33}\ \bar{b}_L b_R + m^d_{23}\ \bar{s}_L b_R + m^d_{13}\ \bar{d}_L b_R + m^d_{12}\ \bar{d}_L s_R + m^d_{22}\ \bar{s}_L s_R + m^d_{32}\ \bar{b}_L s_R \\
&+ m^d_{11}\ \bar{d}_L d_R + m^d_{21}\ \bar{s}_L d_R + m^d_{31}\ \bar{b}_L d_R\Big\} + \text{h.c.}\ , \qquad (16) \\
-\mathcal{L}^u_m &= y_t\frac{\tilde{v}_\Phi}{\sqrt{2}}\bar{t}_L t_R + \frac{Y_u}{\sqrt{2}}\Big\{ \tilde{v}_{G_2}\bar{u}_L c_R + i\tilde{v}_{G_1}\bar{c}_L u_R + \frac{1-i}{2}\bar{t}_L\left(u_R\tilde{v}_{G_2} + c_R\tilde{v}_{G_1}\right)\Big\} \\
&+ \frac{\tilde{v}_\chi}{\sqrt{2}\Lambda}\Big\{ m^t_{33}\ \bar{t}_L t_R + m^t_{23}\ \bar{c}_L t_R + m^t_{13}\ \bar{u}_L t_R + m^t_{12}\ \bar{u}_L c_R + m^t_{22}\ \bar{c}_L c_R + m^t_{32}\ \bar{t}_L c_R \\
&+ m^t_{11}\ \bar{u}_L u_R + m^t_{21}\ \bar{c}_L u_R + m^t_{31}\ \bar{t}_L u_R\Big\} + \text{h.c.}\ , \qquad (17) \\
-\mathcal{L}^\ell_m &= y_\tau\frac{v_\eta}{\sqrt{2}}\bar{\tau}_L\tau_R + \frac{Y_\mu}{\sqrt{2}}\bar{\mu}_L\left(e_R\tilde{v}_{H_2} - \mu_R\tilde{v}_{H_1}\right) \\
&+ \frac{\tilde{v}_\chi}{\sqrt{2}\Lambda}\Big\{ m^\ell_{12}\ \bar{e}_L\mu_R + m^\ell_{22}\ \bar{\mu}_L\mu_R + m^\ell_{11}\ \bar{e}_L e_R + m^\ell_{21}\ \bar{\mu}_L e_R\Big\} + \text{h.c.}\ , \qquad (18)
\end{aligned}$$

with $\tilde{v}_\Psi = v_\Psi e^{i\zeta}, \tilde{v}_\Phi = v_\Phi e^{i\gamma}, \tilde{v}_{G_k} = v_{G_k}e^{i\sigma_k}$, $\tilde{v}_{H_k} = v_{H_k}e^{i\rho_k}$ $(k = 1, 2)$ and $\tilde{v}_\chi = v_\chi e^{i\varphi}$. In the above Eqs. (16), (17) and (18), the entries m^d_{ij}, m^t_{ij} and m^ℓ_{ij} are given in Appendix B. Then, the down-type quark mass matrix $\mathcal{M}_d$ is given by

$$\begin{aligned}
\mathcal{M}_d &= \frac{1}{\sqrt{2}}\begin{pmatrix} iY_d\tilde{v}_{H_1} & 0 & 0 \\ \frac{1-i}{2}Y_d\tilde{v}_{H_2} & \frac{1-i}{2}Y_d\tilde{v}_{H_1} & 0 \\ 0 & Y_d\tilde{v}_{H_2} & y_b\tilde{v}_\Psi \end{pmatrix} + \frac{v_\chi e^{i\varphi}}{\sqrt{2}\Lambda}\begin{pmatrix} m^d_{11} & m^d_{12} & m^d_{13} \\ m^d_{21} & m^d_{22} & m^d_{23} \\ m^d_{31} & m^d_{32} & m^d_{33} \end{pmatrix} \\
&= V^d_L \text{Diag}(m_d, m_s, m_b) V^{d\dagger}_R\ . \qquad (19)
\end{aligned}$$

And, the up-type quark mass matrix $\mathcal{M}_u$ can be explicitly expressed as

$$\begin{aligned}
\mathcal{M}_u &= \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & Y_u\tilde{v}_{G_2} & 0 \\ iY_u\tilde{v}_{G_1} & 0 & 0 \\ \frac{1-i}{2}Y_u\tilde{v}_{G_2} & \frac{1-i}{2}Y_u\tilde{v}_{G_1} & y_t\tilde{v}_\Phi \end{pmatrix} + \frac{v_\chi e^{i\varphi}}{\sqrt{2}\Lambda}\begin{pmatrix} m^t_{11} & m^t_{12} & m^t_{13} \\ m^t_{21} & m^t_{22} & m^t_{23} \\ m^t_{31} & m^t_{32} & m^t_{33} \end{pmatrix} \\
&= V^u_L \text{Diag}(m_u, m_c, m_t) V^{u\dagger}_R\ . \qquad (20)
\end{aligned}$$

Finally, with the VEV alignment in Eq. (6) the charged-lepton mass matrix $\mathcal{M}_\ell$ can be explicitly expressed as

$$\begin{aligned}
\mathcal{M}_\ell &= \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & 0 & 0 \\ -Y_\mu\tilde{v}_{H_2} & Y_\mu\tilde{v}_{H_1} & 0 \\ 0 & 0 & y_\tau v_\eta \end{pmatrix} + \frac{v_\chi e^{i\varphi}}{\sqrt{2}\Lambda}\begin{pmatrix} m^\ell_{11} & m^\ell_{12} & 0 \\ m^\ell_{21} & m^\ell_{22} & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
&= V^\ell_L\ \text{Diag}(m_e, m_\mu, m_\tau)\ V^{\ell\dagger}_R\ . \qquad (21)
\end{aligned}$$

In Eqs. (19), (20) and (21), V_L^f and V_R^f are the diagonalization matrices for $\mathcal{M}_f$.

There exist several empirical fermion mass hierarchies in the up- and down-type quark and charged-lepton sectors calculated from the measured values [8] :

$$\begin{array}{llll} m_u = 2.4 \text{ MeV} & & m_c = 1.27 \text{ GeV} & m_t = 171.2 \text{ GeV} \\ m_d = 4.8 \text{ MeV} & m_s = 104 \text{ MeV} & m_b = 4.2 \text{ GeV} & \\ m_e = 0.511 \text{ MeV} & m_\mu = 105.7 \text{ MeV} & m_\tau = 1.777 \text{ GeV} & \end{array} \tag{22}$$

which implies that the possible quark-lepton symmetry [17] is broken by the masses of quarks and leptons. Thus, it is not expected that the known quark mixing pattern is transmitted to the lepton sector in the exactly same form. In addition, a key point inferred from Eq. (22) is that the mass spectrum of the charged leptons exhibits a similar hierarchical pattern to that of the down-type quarks except for the electron mass which is much smaller than d-quark one, unlike that of the up-type quarks which shows a much stronger hierarchical pattern and top-quark is uniquely biggest. Further, there is another interesting empirical relation

$$|V_{us}| \approx \left(\frac{m_d}{m_s}\right)^{\frac{1}{2}} \approx 3\left(\frac{m_e}{m_\mu}\right)^{\frac{1}{2}} , \tag{23}$$

which has been known for quite a long time [18]. For instance, in terms of the Cabbibo angle $\lambda \equiv \sin\theta_{\rm C} \approx |V_{us}|$, the fermion masses are scaled as $(m_e, m_\mu) \approx (\lambda^5, \lambda^2)\ m_\tau$, $(m_d, m_s) \approx (\lambda^4, \lambda^2)\ m_b$ and $(m_u, m_c) \approx (\lambda^8, \lambda^4)\ m_t$, which may represent the followings: (i) there is at least one Higgs scalar shared by both charged-lepton and down-type quark sectors, or (ii) the mixing matrix of the charged lepton sector is similar to that of the down-type quark sector, and (iii) the CKM matrix is mainly generated by the mixing matrix of the down-type quark sector.

One of most interesting features observed by experiments on the charged fermions is that the mass spectra of quarks and charged leptons are strongly hierarchical, *i.e.*, the masses of third generation fermions are much heavier than those of the first and second generation fermions. For the elements of $\mathcal{M}_f$ given in Eqs. (20) and (19), taking into account the most natural case that the charged fermion masses have the strong hierarchy $m_t \gg m_c \gg m_u$ ($m_{b(\tau)} \gg m_{s(\mu)} \gg m_{d(e)}$) as well as Eq. (22), we make a plausible assumption

$$y_t v_\Phi \gg y_b v_\Psi = y_\tau v_\eta = Y_u v_{G_1} \gg Y_d v_{H_2} \gg Y_u v_{G_2} = Y_d v_{H_1} . \tag{24}$$

Then V_L^f and V_R^f can be determined by diagonalizing the matrices $\mathcal{M}_f \mathcal{M}_f^\dagger$ and $\mathcal{M}_f^\dagger \mathcal{M}_f$, respectively, indicated from Eqs. (21), (20) and (19). Especially, the mixing matrix V_L^f

becomes one of the matrices composing the PMNS and CKM ones and it will be shown later. A general diagonalizing matrix V^f_L can be parameterized in terms of three mixing angles and six phases:

$$V^f_L = \begin{pmatrix} c^f_2 c^f_3 & c^f_2 s^f_3 e^{i\phi^f_3} & s^f_2 e^{i\phi^f_2} \\ -c^f_1 s^f_3 e^{-i\phi^f_3} - s^f_1 s^f_2 c^f_3 e^{i(\phi^f_1 - \phi^f_2)} & c^f_1 c^f_3 - s^f_1 s^f_2 s^f_3 e^{i(\phi^f_1 - \phi^f_2 + \phi^f_3)} & s^f_1 c^f_2 e^{i\phi^f_1} \\ s^f_1 s^f_3 e^{-i(\phi^f_1 + \phi^f_3)} - c^f_1 s^f_2 c^f_3 e^{-i\phi^f_2} & -s^f_1 c^f_3 e^{-i\phi^f_1} - c^f_1 s^f_2 s^f_3 e^{i(\phi^f_3 - \phi^f_2)} & c^f_1 c^f_2 \end{pmatrix} P_f \ , \quad (25)$$

where $s^f_i \equiv \sin\theta^f_i$, $c^f_i \equiv \cos\theta^f_i$ and a diagonal phase matrix $P_f = \text{diag}(e^{i\xi^f_1}, e^{i\xi^f_2}, e^{i\xi^f_3})$ which can be rotated away by the phase redefinition of left-charged fermion fields.

1. *The down-type quark sector and its mixing matrix*

First, we consider the down-type quark sector. From Eq. (19) we see that the down-type quark mass matrix $\mathcal{M}_d$ can be diagonalized in the mass basis by a biunitary transformation, $V^{d\dagger}_L \mathcal{M}_d V^{d(\ell)}_R = \text{Diag}(m_d, m_s, m_b)$. The matrices V^d_L and V^d_R can be determined by diagonalizing the matrices $\mathcal{M}_d \mathcal{M}^\dagger_d$ and $\mathcal{M}^\dagger_d \mathcal{M}_d$, respectively. Especially, the left-handed down-type quark mixing matrix V^d_L becomes one of the matrices composing the CKM matrix such as $V_{\text{CKM}} \equiv V^{u\dagger}_L V^d_L$ [see Eq. (41)]. From Eq. (19) the hermitian square of the mass matrix for down-type quark $\mathcal{M}_d \mathcal{M}^\dagger_d$ can be obtained. And, from Eqs. (19) and (25) the mixing angles and phases can be expressed in terms of Eq. (24) as

$$\begin{aligned} &\theta^d_1 \simeq \frac{v_\chi}{\Lambda}\left(\frac{y^a_b}{2y_b} + \frac{y^s_b}{3y_b}\right) \ , && \theta^d_2 \simeq \frac{v_\chi}{\Lambda}\frac{2y^s_b}{3y_b} \ , && \theta^d_3 \simeq \sqrt{2}\frac{v_{H_1}}{v_{H_2}} \ , \\ &\phi^d_1 \simeq \frac{\varphi}{2} + \frac{\pi}{2} \ , && \phi^d_2 \simeq \frac{\varphi}{2} \ , && \phi^d_3 \simeq \frac{\rho_{12}}{2} + \frac{1}{2}\arg(i-1) \ , \end{aligned} \quad (26)$$

where the parameters y^s_b and y^a_b are positive real numbers of order unity. The empirical relation Eq. (22) for down-type quark can be satisfied by setting as follows

$$\frac{v_\chi}{\Lambda}\left(\frac{y^a_b}{2y_b} + \frac{y^s_b}{3y_b}\right) \equiv A_d \lambda^2 \ , \qquad \frac{v_\chi}{\Lambda}\frac{2y^s_b}{3y_b} \equiv B_d \lambda^3 \ , \qquad v_{H_1} = \frac{\lambda}{\sqrt{2}} v_{H_2} \ , \quad (27)$$

where the parameters A_d and B_d are positive real number of order unity. Note that the third relation in Eq. (27) comes from the renormalizable terms. Then, from the above relation the mass squared eigenvalues are written as

$$\begin{aligned} &V^{d\dagger}_L \mathcal{M}_d \mathcal{M}^\dagger_d V^d_L \equiv \text{Diag}\left(m^2_d, \ m^2_s, \ m^2_b\right) \\ &\simeq \text{Diag}\left(\frac{\lambda v^2_{H_2}}{\sqrt{2}} Y^2_d \frac{v_\chi}{\Lambda}\left\{Y_1 \sin(\rho_{12} - \varphi) + Y_2 \cos(\rho_{12} - \varphi)\right\}, \ \frac{1}{4} Y^2_d v^2_{H_2}, \ \frac{1}{2} y^2_b v^2_\Psi\right) \ , \end{aligned} \quad (28)$$

where $Y_d v_{H_2} = \sqrt{2}\lambda^2 y_b v_\Psi$, and $Y_1 = \frac{Y_1^d}{Y_d} + \frac{Y_d^a}{4Y_d} + \frac{Y_d^s}{6Y_d}$, $Y_2 = \frac{Y_d^a}{4Y_d} + \frac{Y_d^s}{6Y_d}$. So, one can obtain the measured value of $m_d/m_b \simeq \lambda^4$. Then, we can obtain the mixing matrix V_L^d of the down-type quarks: under the constraint of unitarity up to $\mathcal{O}(\lambda^3)$, it can be written as

$$V_L^d = \begin{pmatrix} 1-\frac{\lambda^2}{2} & \lambda e^{i\phi_3^d} & B_d\lambda^3 e^{i\frac{\varphi}{2}} \\ -\lambda e^{-i\phi_3^d} & 1-\frac{\lambda^2}{2} & A_d\lambda^2 e^{i(\frac{\varphi}{2}+\frac{\pi}{2})} \\ A_d\lambda^3 e^{-i(\phi_3^d+\frac{\varphi}{2}+\frac{\pi}{2})} - B_d\lambda^3 e^{-i\frac{\varphi}{2}} & -A_d\lambda^2 e^{-i(\frac{\varphi}{2}+\frac{\pi}{2})} & 1 \end{pmatrix} P_d + \mathcal{O}(\lambda^4)\ . \quad (29)$$

2. The up-type quark sector and its mixing matrix

Next, let us consider the up-type quark sector to obtain the realistic CKM matrix. From Eq. (20) and Eq. (24) the hermitian square of the mass matrix for up-type quark $\mathcal{M}_u\mathcal{M}_u^\dagger$, with the condition given in Eq. (24) the mass squared eigenvalues are written in a good approximation as

$$m_t^2 \simeq \frac{1}{2}y_t^2 v_\Phi^2\ , \qquad m_c^2 \simeq \frac{1}{2}Y_u^2 v_{G_1}^2\ , \qquad m_u^2 \simeq \frac{1}{2}Y_u^2 v_{G_2}^2\ . \quad (30)$$

Then, one can set $m_u/m_t \simeq v_{G_2}/v_\Phi \simeq \lambda^8$ and $m_c/m_t \simeq v_{G_1}/v_\Phi \simeq \lambda^4$ for equal amounts of Yukawa couplings. And, the mixing angles and phases can be expressed in terms of Eq. (24) as

$$\begin{aligned} &\theta_1^u \simeq \frac{v_\chi}{\Lambda}\left(\frac{y_t^a}{2y_t}+\frac{y_t^s}{3y_t}\right)\ , \qquad \theta_2^u \simeq \frac{v_\chi}{\Lambda}\frac{2y_t^s}{3y_t}\ , \qquad \theta_3^u \simeq \frac{v_\chi}{\Lambda}\left(\frac{Y_u^a}{2Y_u}+\frac{Y_u^s}{3Y_u}\right) \\ &\phi_1^u \simeq \frac{\varphi}{2}+\frac{\pi}{2}\ , \qquad \phi_2^u \simeq \frac{\varphi}{2}\ , \qquad \phi_3^u \simeq \frac{\pi}{2}+\frac{\varphi}{2}\ . \end{aligned} \quad (31)$$

Due to $v_\chi/\Lambda \sim \lambda^2$ in Eq. (27) and the measured value of $m_u/m_c \approx v_{G_2}/v_{G_1} \approx \lambda^4$ in Eq. (22), it is impossible to generate the Cabbibo angle, $\lambda \approx |V_{us}|$, from the mixing between the first and second generations in the up-type quark sector: if one sets $|(V_L^u)_{12}| = \theta_3^u$, then from Eqs. (30) and (31) one obtains $|(V_L^u)_{12}| \sim v_\chi/\Lambda \approx \lambda^2$, in discrepancy with the measured $\lambda \approx |V_{us}|$. And similar to Eq. (27), from Eq. (31) one can set

$$\theta_1^u \simeq A_u\lambda^2\ , \qquad \theta_2^u \simeq B_u\lambda^3\ , \qquad \theta_3^u \simeq C_u\lambda^2\ , \quad (32)$$

where A_u, B_u, C_u are positive real numbers of order unity. Then, plugging Eqs. (31) and (32) into Eq. (25) the up-type quark mixing matrix V_L^u can be written, under the constraint of unitarity up to $\mathcal{O}(\lambda^3)$, as

$$V_L^u = \begin{pmatrix} 1 & C_u\lambda^2 e^{i\frac{\varphi+\pi}{2}} & B_u\lambda^3 e^{i\frac{\varphi}{2}} \\ -C_u\lambda^2 e^{-i\frac{\varphi+\pi}{2}} & 1 & A_u\lambda^2 e^{i(\frac{\varphi}{2}+\frac{\pi}{2})} \\ -B_u\lambda^3 e^{-i\frac{\varphi}{2}} & -A_u\lambda^2 e^{-i(\frac{\varphi}{2}+\frac{\pi}{2})} & 1 \end{pmatrix} P_u + \mathcal{O}(\lambda^4) \ , \tag{33}$$

which indicates that the mixing matrix in the up-type quark sector can affect, at most, the next leading order contributions in λ. Flavor-changing neutral currents (FCNCs) in up-type quarks may restrict $A_u, B_u, C_u < \lambda$, which we will not discuss.

3. *The charged lepton sector and its mixing matrix*

Similar to the quark sector, from Eq. (21) we see that the charged lepton mass matrix $\mathcal{M}_\ell$ can be diagonalized in the mass basis by a biunitary transformation, $V_L^{\ell\dagger}\mathcal{M}_\ell V_R^{d(\ell)} = \text{Diag}(m_e, m_\mu, m_\tau)$. The matrices V_L^ℓ and V_R^ℓ can be determined by diagonalizing the matrices $\mathcal{M}_\ell\mathcal{M}_\ell^\dagger$ and $\mathcal{M}_\ell^\dagger\mathcal{M}_\ell$, respectively. Especially, the charged lepton mixing matrix V_L^ℓ becomes one of the matrices composing the PMNS matrix such as $U_{\text{PMNS}} \equiv V_L^{\ell\dagger}U_\nu$, respectively [see Eq. (60)]. From Eq. (21) the hermitian square of the mass matrix for charged lepton $\mathcal{M}_\ell\mathcal{M}_\ell^\dagger$ can be obtained. From Eqs. (21) and (25) the mixing angles and phases can be expressed in terms of Eq. (24) as

$$\theta_3^\ell \simeq \frac{1}{\sqrt{2}}\frac{Y_2^\mu}{Y_\mu}\frac{v_\chi}{\Lambda} \ , \qquad\qquad \phi_3^\ell \simeq \frac{\varphi}{2} + \frac{1}{2}\arg(i-1) \ , \tag{34}$$

and $\phi_1^\ell = \phi_2^\ell = 0, \theta_1^\ell = \theta_2^\ell = 0$, where the parameters Y_2^μ, Y_μ are positive real numbers of order unity. Then, from the above relation the mass squared eigenvalues are written, in a good approximation, as

$$V_L^{\ell\dagger}\mathcal{M}_\ell\mathcal{M}_\ell^\dagger V_L^{d(\ell)} \equiv \text{Diag}\left(m_e^2,\ m_\mu^2,\ m_\tau^2\right) \simeq \text{Diag}\left(\frac{1}{2}Y_2^{\mu 2}v_{H_2}^2\left(\frac{v_\chi}{\Lambda}\right)^2,\ \frac{1}{2}Y_\mu^2 v_{H_2}^2,\ \frac{1}{2}y_\tau^2 v_\eta^2\right) . \tag{35}$$

Due to the ratio of the measured values $m_e/m_\mu \simeq \lambda^3$ and $m_\mu/m_\tau \simeq \lambda^2$, they can be expressed in terms of Eq. (21) as $m_e/m_\mu \simeq Y_2^\mu v_\chi/(Y_\mu\Lambda)$ and $m_\mu/m_\tau \simeq Y_\mu v_{H_2}/(y_\tau v_\eta)$. And, one can express

$$\frac{v_\chi}{\Lambda}\frac{Y_2^\mu}{\sqrt{2}Y_\mu} = A_\ell\lambda^3 \ , \tag{36}$$

where the parameter A_ℓ is positive real number of order unity. Then, we can obtain the mixing matrix V_L^ℓ of the charged leptons: under the constraint of unitarity up to $\mathcal{O}(\lambda^3)$, it can be written as

$$V_L^\ell = \begin{pmatrix} 1 & A_\ell \lambda^3 e^{i\phi_3^\ell} & 0 \\ -A_\ell \lambda^3 e^{-i\phi_3^\ell} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} P_\ell + \mathcal{O}(\lambda^4)\ . \tag{37}$$

It indicates that the effect of mixing in the charged-lepton sector to the PMNS matrix is at least less than $\lambda^3 \approx 0.2^\circ$, and its contribution to the PMNS matrix is negligible because of the relatively large reactor angle θ_{13} measured in Daya Bay and RENO experiments [9].

III. PMNS AND CKM MIXING MATRICES AND MASS SPECTRA

A. Quark Sector

In the weak eigenstate basis, the mass terms in Eqs. (16) and (17) and the charged gauge interactions can be written as

$$-\mathcal{L}^q_{mW} = \overline{q^u_L}\mathcal{M}_u q^u_R + \overline{q^d_L}\mathcal{M}_d q^d_R + \frac{g}{\sqrt{2}} W^+_\mu\ \overline{q^u_L}\gamma^\mu q^d_L + \text{h.c.}\ . \tag{38}$$

Let us first consider the quark sector. From Eq. (38), to diagonalize the up- and down-type quark mass matrices such that

$$V_L^{f\dagger}\mathcal{M}_f V_R^f = \text{Diag}(m_{f_1}, m_{f_2}, m_{f_3})\ , \tag{39}$$

we can rotate the quark fields from the weak eigenstates to the mass eigenstates:

$$q_L^{u(d)} \to V_L^{u(d)\dagger} q_L^{u(d)}\ , \qquad q_R^{u(d)} \to V_R^{u(d)\dagger} q_R^{u(d)}\ . \tag{40}$$

Then, from the charged current terms in Eq. (38), we obtain the CKM matrix

$$V_{\text{CKM}} = V_L^{u\dagger} V_L^d\ . \tag{41}$$

From Eqs. (29) and (33) we can obtain directly $V_L^{u\dagger}V_L^d$, and by recasting the result with the transformations $d \to de^{i\xi'}$, $c \to ce^{i(\phi_3^d - \xi + \xi')}$, $s \to se^{i(\phi_3^d - \xi)}$, $t \to te^{i(\phi_3^d + \frac{\varphi}{2} + \frac{\pi}{2} - \xi)}$ and

$b \to b e^{i(\phi_3^d+\frac{\varphi}{2}+\frac{\pi}{2}-\xi)}$, we can rewrite the CKM matrix as [4]

$$\begin{aligned} V_{\rm CKM} &= \begin{pmatrix} 1-\frac{\lambda^2}{2}+C_u\lambda^3\sin(\phi_3^d-\frac{\varphi}{2}) & \lambda - C_u\lambda^2\sin(\phi_3^d-\frac{\varphi}{2}) & B\lambda^3 e^{-i(\phi_3^d+\frac{\pi}{2}-\xi)} \\ -\lambda + C_u\lambda^2\sin(\phi_3^d-\frac{\varphi}{2}) & 1-\frac{\lambda^2}{2}+C_u\lambda^3\sin(\phi_3^d-\frac{\varphi}{2}) & A\lambda^2 e^{-i\xi'} \\ (A-Be^{i(\phi_3^d+\frac{\pi}{2}-\xi)})\lambda^3 e^{-i\xi'} & -A\lambda^2 & 1 \end{pmatrix} \\ &+ \mathcal{O}(\lambda^4) \ . \end{aligned} \tag{42}$$

where [5] $A = A_d - A_u$, $B = B_d - B_u$, $\xi' \simeq C_u\lambda^3\cos(\frac{\varphi}{2}-\phi_3^d)$ and $\xi \simeq C_u\lambda\cos(\frac{\varphi}{2}-\phi_3^d)$. If one set $C_u \to 0$ which can be realized by $C_u \sim \frac{Y_u^a}{2Y_u}+\frac{Y_u^s}{3Y_u} \simeq \lambda^n$ with $n \geq 1$, and

$$B e^{-i(\phi_3^d+\frac{\pi}{2}-\xi)} = A(\rho + i\eta) \ , \tag{43}$$

then one can obtain the realistic CKM matrix in the Wolfenstein parametrization [19] given by

$$V_{\rm CKM} = \begin{pmatrix} 1-\frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho+i\eta) \\ -\lambda & 1-\frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1-\rho+i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4) \ . \tag{44}$$

As reported in Ref. [12] the best-fit values of the parameters λ, A, $\bar\rho$, $\bar\eta$ with 1σ errors are

$$\begin{aligned} \lambda &= \sin\theta_C = 0.22543 \pm 0.00077 \ , \qquad A = 0.812^{+0.013}_{-0.027} \ , \\ \bar\rho &= 0.144 \pm 0.025 \ , \qquad\qquad\qquad\qquad \bar\eta = 0.342^{+0.016}_{-0.015} \ , \end{aligned} \tag{45}$$

where $\bar\rho = \rho(1-\lambda^2/2)$ and $\bar\eta = \eta(1-\lambda^2/2)$. The effects caused by CP violation are always proportional to the Jarlskog invariant [20], defined as $J_{CP}^{\rm quark} = {\rm Im}[V_{ud}V_{cs}V^*_{us}V^*_{cd}] \simeq A^2\lambda^6\eta$ whose value is $2.96^{+0.18}_{-0.17}\times 10^{-5}$ at 1σ level [12].

From Eqs. (28) and (30), the observed quark masses respect the following relation

$$\begin{aligned} m_d : \ m_s : \ m_b &\simeq v_{H_2}Y_d\sqrt{\frac{\lambda v_\chi}{\sqrt{2}\Lambda}\left\{Y_1\sin(\rho_{12}-\varphi)+Y_2\cos(\rho_{12}-\varphi)\right\}} : \ \frac{1}{2}Y_d v_{H_2} : \ \frac{1}{\sqrt{2}}y_b v_\Psi \ , \\ m_u : \ m_c : \ m_t &\simeq \frac{1}{\sqrt{2}}Y_u v_{G_2} : \ \frac{1}{\sqrt{2}}Y_u v_{G_1} : \ \frac{1}{\sqrt{2}}y_t v_\Phi \ . \end{aligned} \tag{46}$$

---

[4] Note here that since the matrix in Eq. (42) has the phase φ dependence which is from χ field, when $C_u \to \lambda^n$ with $n \geq 1$ (n: integer) one can re-parameterize and obtain the CKM matrix.

[5] Taking into account the FCNCs in the up-type quarks we may approximate $A_u, B_u, C_u \to 0$ [see below Eq. (33)].

B. Lepton Sector

The mass terms in Eqs. (8) and (18) and the charged gauge interactions in the weak eigenstate basis can be written in (block) matrix form as, using $\overline{N^c_R} m_D \nu^c_L = \overline{\nu_L} m_D^T N_R$,

$$\begin{aligned} -\mathcal{L}^\ell_{mW} &= \frac{1}{2}\overline{N^c_R} M_R N_R + \overline{\nu_L} m_D N_R + \overline{\ell_L}\mathcal{M}_\ell \ell_R + \frac{g}{\sqrt{2}} W^-_\mu \overline{\ell_L}\gamma^\mu \nu_L + \text{h.c.} \qquad (47) \\ &= \frac{1}{2}\begin{pmatrix} \overline{\nu_L} & \overline{N^c_R} \end{pmatrix} \begin{pmatrix} 0 & m_D \\ m_D^T & M_R \end{pmatrix} \begin{pmatrix} \nu^c_L \\ N_R \end{pmatrix} + \overline{\ell_L}\mathcal{M}_\ell \ell_R + \frac{g}{\sqrt{2}} W^-_\mu \overline{\ell_L}\gamma^\mu \nu_L + \text{h.c.} \qquad (48) \end{aligned}$$

Here $\ell = (e, \mu, \tau)$, $\nu = (\nu_e, \nu_\mu, \nu_\tau)$, $N_R = (N_{R1}, N_{R2}, N_{R3})$.

To find the neutrino masses and mixing matrix we are to diagonalize the 6×6 matrix

$$\begin{pmatrix} 0 & m_D \\ m_D^T & M_R \end{pmatrix} . \tag{49}$$

We start by diagonalizing M_R. For this purpose, we perform a basis rotation $\widehat{N}_R = U_R^\dagger N_R$, so that the right-handed Majorana mass matrix M_R becomes a diagonal matrix $\widehat{M_R}$ with real and positive mass eigenvalues $M_1 = aM$, $M_2 = M$ and $M_3 = bM$,

$$\widehat{M_R} = U_R^T M_R U_R = M\ U_R^T \begin{pmatrix} 1 + \frac{2}{3}\kappa e^{i\varphi} & -\frac{1}{3}\kappa e^{i\varphi} & -\frac{1}{3}\kappa e^{i\varphi} \\ -\frac{1}{3}\kappa e^{i\varphi} & \frac{2}{3}\kappa e^{i\varphi} & 1 - \frac{1}{3}\kappa e^{i\varphi} \\ -\frac{1}{3}\kappa e^{i\varphi} & 1 - \frac{1}{3}\kappa e^{i\varphi} & \frac{2}{3}\kappa e^{i\varphi} \end{pmatrix} U_R = \begin{pmatrix} aM & 0 & 0 \\ 0 & M & 0 \\ 0 & 0 & bM \end{pmatrix}, \tag{50}$$

where $\kappa = y^\nu_R v_\chi / M$. We find $a = \sqrt{1 + \kappa^2 + 2\kappa\cos\varphi}$, $b = \sqrt{1 + \kappa^2 - 2\kappa\cos\varphi}$, and a diagonalizing matrix

$$U_R = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} e^{i\frac{\psi_1}{2}} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{i\frac{\psi_2}{2}} \end{pmatrix} , \tag{51}$$

with phases

$$\psi_1 = \tan^{-1}\left(\frac{-\kappa\sin\varphi}{1 + \kappa\cos\varphi}\right) \quad \text{and} \quad \psi_2 = \tan^{-1}\left(\frac{\kappa\sin\varphi}{1 - \kappa\cos\varphi}\right) . \tag{52}$$

As the magnitude of κ defined in Eq. (50) decreases, the phases $\psi_{1,2}$ go to 0 or π. With the basis rotation $N_R \to U_R^\dagger N_R$, the Dirac neutrino mass matrix gets modified to

$$m_D \to \widetilde{m}_D = m_D U_R = \frac{v_\Phi e^{i\gamma}}{\sqrt{2}} y^\nu_3 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & y_2 \\ 0 & y_1 & 0 \end{pmatrix} U_R\ , \tag{53}$$

where $y_1 = y^\nu_1/y^\nu_3, y_2 = y^\nu_2/y^\nu_3$. At this point,

$$-\mathcal{L}_{mW} = \frac{1}{2}\begin{pmatrix}\overline{\nu_L} & \overline{\widehat{N}^c_R}\end{pmatrix}\begin{pmatrix}0 & \widetilde{m}_D \\ \widetilde{m}^T_D & \widehat{M}_R\end{pmatrix}\begin{pmatrix}\nu^c_L \\ \widehat{N}_R\end{pmatrix} + \overline{\ell_L}\mathcal{M}_\ell\ell_R + \frac{g}{\sqrt{2}}W^-_\mu\overline{\ell_L}\gamma^\mu\nu_L + \text{h.c.}\ . \tag{54}$$

Now we take the limit of large M (seesaw mechanism) and focus on the mass matrix of the light neutrinos M_ν,

$$-\mathcal{L}_{mW} = \frac{1}{2}\overline{\nu_L}\mathcal{M}_\nu\nu^c_L + \overline{\ell_L}\mathcal{M}_\ell\ell_R + \frac{g}{\sqrt{2}}W^-_\mu\overline{\ell_L}\gamma^\mu\nu_L + \text{h.c.} + \text{terms in } N_R \tag{55}$$

with

$$\mathcal{M}_\nu = -\widetilde{m}_D\,\widehat{M}^{-1}_R\,\widetilde{m}^T_D. \tag{56}$$

We perform basis rotations from weak to mass eigenstates in the leptonic sector,

$$\widehat{\ell}_L = V^{\ell\dagger}_L\ell_L\ , \qquad \widehat{\ell}_R = V^{\ell\dagger}_R\ell_R\ , \qquad \widehat{\nu}_L = U^\dagger_\nu\nu_L\ , \tag{57}$$

where $U_\nu, V_{L(R)}$ are unitary matrices chosen so as the matrices

$$\begin{aligned}\widehat{\mathcal{M}}_\nu &= U^\dagger_\nu\mathcal{M}_\nu U^*_\nu = -U^\dagger_\nu m_D U_R\widehat{M}^{-1}_R(U^\dagger_\nu m_D U_R)^T\ , \\ \widehat{\mathcal{M}}_\ell &= V^{\ell\dagger}_L\mathcal{M}_\ell V_R\end{aligned} \tag{58}$$

are diagonal. From Eqs. (35) and (36) the observed charged lepton masses respect

$$m_e\ :\ m_\mu\ :\ m_\tau \simeq A_\ell\lambda^3 Y_\mu v_{H_2}\ :\ \frac{Y_\mu}{\sqrt{2}}v_{H_2}\ :\ \frac{y_\tau}{\sqrt{2}}v_\eta\ . \tag{59}$$

And from the charged current term in Eq. (54) we obtain the lepton mixing matrix U_{PMNS} as

$$U_{\text{PMNS}} = V^{\ell\dagger}_L U_\nu. \tag{60}$$

The matrix U_{PMNS} can be written in terms of three mixing angles and three CP-odd phases (one for the Dirac neutrinos and two for the Majorana neutrinos) as [8]

$$U_{\text{PMNS}} = \begin{pmatrix} c_{13}c_{12} & c_{13}s_{12} & s_{13}e^{-i\delta_{CP}} \\ -c_{23}s_{12} - s_{23}c_{12}s_{13}e^{i\delta_{CP}} & c_{23}c_{12} - s_{23}s_{12}s_{13}e^{i\delta_{CP}} & s_{23}c_{13} \\ s_{23}s_{12} - c_{23}c_{12}s_{13}e^{i\delta_{CP}} & -s_{23}c_{12} - c_{23}s_{12}s_{13}e^{i\delta_{CP}} & c_{23}c_{13}\end{pmatrix}P_\nu\ , \tag{61}$$

where $P_\nu = \text{Diag}(e^{-i\varphi_1/2}, e^{-i\varphi_2/2}, 1)$, and $s_{ij} \equiv \sin\theta_{ij}$ and $c_{ij} \equiv \cos\theta_{ij}$.

After seesawing, the light neutrino mass matrix is given by [6]

$$\begin{aligned}\mathcal{M}_\nu &= -\widetilde{m}_D \widehat{M}_R^{-1} \widetilde{m}_D^T \\ &= e^{2i\gamma} m_0 \begin{pmatrix} 1+\frac{2e^{i\psi_1}}{a} & (1-\frac{e^{i\psi_1}}{a})y_2 & (1-\frac{e^{i\psi_1}}{a})y_1 \\ (1-\frac{e^{i\psi_1}}{a})y_2 & (1+\frac{e^{i\psi_1}}{2a}+\frac{3e^{i\psi_2}}{2b})y_2^2 & (1+\frac{e^{i\psi_1}}{2a}-\frac{3e^{i\psi_2}}{2b})y_1 y_2 \\ (1-\frac{e^{i\psi_1}}{a})y_1 & (1+\frac{e^{i\psi_1}}{2a}-\frac{3e^{i\psi_2}}{2b})y_1 y_2 & (1+\frac{e^{i\psi_1}}{2a}+\frac{3e^{i\psi_2}}{2b})y_1^2 \end{pmatrix} ,\end{aligned} \tag{62}$$

where we have defined an overall scale $m_0 = v_\Phi^2 y_3^{\nu 2}/(6M)$ for the light neutrino masses. And the overall phase can be rotated away by redefining the light neutrino field. The mass matrix $\mathcal{M}_\nu$ is diagonalized by the mixing matrix U_ν,

$$\mathcal{M}_\nu = U_\nu \ \mathrm{Diag}(m_{\nu_1}, m_{\nu_2}, m_{\nu_3}) \ U_\nu^T . \tag{63}$$

Here m_{ν_i} $(i = 1, 2, 3)$ are the light neutrino masses. Interestingly, the mixing matrix U_R in Eq. (51) reflects an exact TBM. Therefore Eq. (62) directly indicates that there could be deviations from the exact TBM if the Dirac neutrino Yukawa couplings do not have the same magnitude. In the limit $y_1^\nu = y_2^\nu$, the mass matrix in Eq. (62) acquires a μ–τ symmetry [21] that leads to $\theta_{13}^\nu = 0$ and $\theta_{23}^\nu = -\pi/4$. Moreover, in the limit $y_1^\nu = y_2^\nu = y_3^\nu$ $(y_1, y_2 \to 1)$, the mass matrix (62) gives the TBM angles and the corresponding mass eigenvalues:

$$\begin{aligned} \theta_1^\nu &= -\frac{\pi}{4} , \qquad \theta_2^\nu = 0 , \qquad \theta_3^\nu = \sin^{-1}\left(\frac{1}{\sqrt{3}}\right) , \\ m_{\nu_1} &= \frac{3m_0}{a} , \qquad m_{\nu_2} = 3m_0 , \qquad m_{\nu_3} = \frac{3m_0}{b} . \end{aligned} \tag{64}$$

These mass eigenvalues are disconnected from the mixing angles. The neutrino texture in Eq. (62) provides naturally the mildness of neutrino masses, because the components giving the TBM are multiplied and constrained by neutrino Yukawa couplings.

Due to in general $y_1, y_2 \neq 1$, there are deviations from their TBM values. Moreover, recent neutrino data, *i.e.* $\theta_{13} \neq 0$, require deviations of $y_{1,2}$ from unity because the contribution of V_L^ℓ in Eq. (37) from the charged-lepton sector is expected to be small, leading to a possibility to search for CP violation in neutrino oscillation experiments. To diagonalize the above mass matrix Eq. (62), we consider the hermitian matrix $\mathcal{M}_\nu \mathcal{M}_\nu^\dagger$, from which we obtain the masses and mixing angles:

---

[6] The neutrino mass matrix form given by Eq. (62) is different from the one given in Ref. [5] due to the opposite sign in front of $\frac{3e^{i\psi_2}}{2b}$ in 2-3 sector of the light neutrino mass matrix. So, it makes a difference in numerical results.

$$\mathcal{M}_\nu \mathcal{M}_\nu^\dagger = \begin{pmatrix} A & B & C \\ B^* & F & |G|e^{i\phi_1^\nu} \\ C^* & |G|e^{-i\phi_1^\nu} & K \end{pmatrix} = U_\nu \ \mathrm{Diag}(m_{\nu_1}^2, m_{\nu_2}^2, m_{\nu_3}^2) \ U_\nu^\dagger \ , \tag{65}$$

where

$$\begin{aligned} A &= p+q+2g_1 \ , \quad F = y_2^2\left(p+\frac{q}{4}+r-g_1-g_2\right) \ , \quad K = y_1^2\left(p+\frac{q}{4}+r-g_1+g_2\right) \ , \\ B &= y_2\left(p-\frac{q}{2}+\frac{g_1}{2}-g_3+i\frac{3(g_4+g_5)}{2a}\right) \ , \ C = y_1\left(p-\frac{q}{2}+\frac{g_1}{2}+g_3+i\frac{3(g_4-g_5)}{2a}\right) \ , \\ G &= y_1 y_2\left(p+\frac{q}{4}-r-g_1+ig_6\right) \ , \end{aligned} \tag{66}$$

with

$$\begin{aligned} p &= m_0^2(1+y_1^2+y_2^2) \ , \qquad q = m_0^2\frac{4+y_1^2+y_2^2}{a^2} \ , \qquad r = 9m_0^2\frac{y_1^2+y_2^2}{4b^2} \ , \\ g_1 &= m_0^2\cos\psi_1\frac{2-y_1^2-y_2^2}{a} \ , \qquad g_2 = 3m_0^2(y_1^2-y_2^2)\frac{\cos\gamma+2a\cos\psi_2}{2ab} \ , \\ g_3 &= 3m_0^2(y_1^2-y_2^2)\frac{a\cos\psi_2-\cos\gamma}{2ab} \ , \qquad g_4 = m_0^2(2-y_1^2-y_2^2)\sin\psi_1 \ , \\ g_5 &= m_0^2(y_1^2-y_2^2)\frac{a\sin\psi_2+\sin\gamma}{b} \ , \qquad g_6 = 3m_0^2(y_1^2-y_2^2)\frac{\sin\gamma-2a\sin\psi_2+}{2ab} \ , \end{aligned} \tag{67}$$

and $\gamma \equiv \psi_1 - \psi_2 = \tan^{-1}\left(\frac{2\kappa\sin\varphi}{\kappa^2-1}\right)$. In the limit of $y_1, y_2 \to 1$ the parameters in Eq. (67) behave as $p \to 3m_0^2, q \to 6m_0^2/a^2, r \to 2m_0^2/b^2$ and $g_i \to 0$. So, as expected, the matrix in Eq. (65) gives the TBM values. Similarly Eq. (25), we have three mixing angles $(\theta_1^\nu, \theta_2^\nu, \theta_3^\nu)$, three phases $(\phi_1^\nu, \phi_2^\nu, \phi_3^\nu)$, and the three mass-squared eigenvalues. In turn, this mixing matrix U_ν becomes one of the mixing matrix composing the PMNS matrix. To see how the neutrino mass matrix given by Eq.(62) can lead to deviations from their TBM values, we first introduce three small quantities $\varepsilon_i, \ (i=1,2,3)$, which are responsible for the deviations of the θ_j from their TBM values:

$$\theta_1^\nu = -\frac{\pi}{4}+\varepsilon_1 \ , \qquad \theta_2^\nu = \varepsilon_2 \ , \qquad \theta_3^\nu = \sin^{-1}\left(\frac{1}{\sqrt{3}}\right)+\varepsilon_3 \ . \tag{68}$$

Then the mixing matrix U_ν up to order ε_i can be written as

$$\begin{aligned} U_\nu \ = & \begin{pmatrix} \frac{\sqrt{2}-\varepsilon_3}{\sqrt{3}} & \frac{1+\varepsilon_3\sqrt{2}}{\sqrt{3}}e^{i\phi_3^\nu} & \varepsilon_2 e^{i\phi_2^\nu} \\ -\frac{(1+\varepsilon_1+\varepsilon_3\sqrt{2})e^{-i\phi_3^\nu}}{\sqrt{6}}+\frac{\varepsilon_2 e^{i(\phi_1^\nu-\phi_2^\nu)}}{\sqrt{3}} & \frac{\sqrt{2}+\varepsilon_1\sqrt{2}-\varepsilon_3}{\sqrt{6}}+\frac{\varepsilon_2 e^{i(\phi_1^\nu-\phi_2^\nu+\phi_3^\nu)}}{\sqrt{6}} & \frac{(-1+\varepsilon_1)e^{i\phi_1^\nu}}{\sqrt{2}} \\ -\frac{(1-\varepsilon_1-\varepsilon_3\sqrt{2})e^{-i(\phi_1^\nu+\phi_3^\nu)}}{\sqrt{6}}-\frac{\varepsilon_2 e^{-i\phi_2^\nu}}{\sqrt{3}} & \frac{(\sqrt{2}-\varepsilon_3-\sqrt{2}\varepsilon_1)e^{-i\phi_1^\nu}}{\sqrt{6}}-\frac{\varepsilon_2 e^{i(\phi_3^\nu-\phi_2^\nu)}}{\sqrt{6}} & \frac{1+\varepsilon_1}{\sqrt{2}} \end{pmatrix} P_\nu \\ & + \ \mathcal{O}(\varepsilon_i^2) \ . \end{aligned} \tag{69}$$

Now, the straightforward calculation with the general parametrization of U_ν in Eq. (25) leads to the expressions for the masses and mixing parameters

$$\begin{aligned}
\tan\theta_1^\nu &= \frac{\mathrm{Im}[C]\sin\phi_2^\nu - \mathrm{Re}[C]\cos\phi_2^\nu}{\mathrm{Im}[B]\cos(\phi_1^\nu-\phi_2^\nu)+\mathrm{Re}[B]\sin(\phi_1^\nu-\phi_2^\nu)}\ , \qquad \phi_1^\nu = \arg(G)\ ,\\
\tan 2\theta_2^\nu &= 2\frac{|c_1^\nu C + e^{i\phi_1^\nu}s_1^\nu B|}{\lambda_3 - A}\ , \qquad \phi_2^\nu = \arg\left(c_1^\nu C + e^{i\phi_1^\nu}s_1^\nu B\right)\ ,\\
\tan 2\theta_3^\nu &= 2\frac{|Z|}{\lambda_2-\lambda_1}\ , \qquad \phi_3^\nu = \arg(Z)\ , \qquad (70)
\end{aligned}$$

where $c_i^\nu = \cos\theta_i^\nu$, $s_i^\nu = \sin\theta_i^\nu$, and

$$\begin{aligned}
\lambda_1 &= A c_2^{\nu 2} - |c_1^\nu C + e^{i\phi_1^\nu}s_1^\nu B|\sin 2\theta_2^\nu + \lambda_3 s_2^{\nu 2}\ ,\\
\lambda_2 &= F c_1^{\nu 2} - |G|\sin 2\theta_1^\nu + K s_1^{\nu 2}\ , \qquad \lambda_3 = K c_1^{\nu 2} + |G|\sin 2\theta_1^\nu + F s_1^{\nu 2}\ ,\\
Z &= c_2^\nu(c_1^\nu B - e^{-i\phi_1^\nu}s_1^\nu C) + s_2^\nu e^{i(\phi_2^\nu-\phi_1^\nu)}\left(\sin 2\theta_1^\nu\frac{K-F}{2} - |G|\cos 2\theta_1^\nu\right)\ . \qquad (71)
\end{aligned}$$

And the squared-mass eigenvalues are given by

$$m_{\nu_1}^2 = \frac{\lambda_1 c_3^{\nu 2} - \lambda_2 s_3^{\nu 2}}{\cos 2\theta_3^\nu}\ , \quad m_{\nu_2}^2 = \frac{\lambda_2 c_3^{\nu 2} - \lambda_1 s_3^{\nu 2}}{\cos 2\theta_3^\nu}\ , \quad m_{\nu_3}^2 = \lambda_3 + |c_1^\nu C + e^{i\phi_1^\nu}s_1^\nu B|\tan\theta_2^\nu\ . \qquad (72)$$

As is well-known, because of the observed hierarchy $\Delta m^2_{\mathrm{Atm}} \gg \Delta m^2_{\mathrm{Sol}} \equiv m_2^2 - m_1^2 > 0$, and the requirement of a Mikheyev-Smirnov-Wolfenstein resonance for solar neutrinos, there are two possible neutrino mass spectra: (i) the normal mass hierarchy (NMH) $m_1 < m_2 < m_3$, and (ii) the inverted mass hierarchy (IMH) $m_3 < m_1 < m_2$. The solar and atmospheric mass-squared differences are given by

$$\Delta m^2_{\mathrm{sol}} \equiv m_{\nu_2}^2 - m_{\nu_1}^2 = \frac{2\,|Z|}{\sin 2\theta_3^\nu}\ , \qquad \Delta m^2_{\mathrm{atm}} \equiv \begin{cases} m_{\nu_3}^2 - m_{\nu_1}^2, & \text{for NMH}\\ m_{\nu_2}^2 - m_{\nu_3}^2, & \text{for IMH}\end{cases}\ , \qquad (73)$$

which are constrained by the neutrino oscillation experimental results. We will discuss it numerically in the next section.

Plugging Eqs. (37) and (69) into Eq. (60), the PMNS matrix is recast to

$$U_{\mathrm{PMNS}} = \begin{pmatrix} U_{\nu 11} - A_\ell\lambda^3 e^{i\phi_3^\ell}U_{\nu 21} & U_{\nu 12} - A_\ell\lambda^3 e^{i\phi_3^\ell}U_{\nu 22} & U_{\nu 13} - A_\ell\lambda^3 e^{i\phi_3^\ell}U_{\nu 23}\\ U_{\nu 21} + A_\ell\lambda^3 e^{-i\phi_3^\ell}U_{\nu 11} & U_{\nu 22} + A_\ell\lambda^3 e^{-i\phi_3^\ell}U_{\nu 12} & U_{\nu 23} + A_\ell\lambda^3 e^{-i\phi_3^\ell}U_{\nu 13}\\ U_{\nu 31} & U_{\nu 32} & U_{\nu 33}\end{pmatrix} P_\nu\ , \qquad (74)$$

where the phase ϕ_3^ℓ is given as Eq. (34). From Eq. (74), the neutrino mixing parameters can be displayed in terms of the standard parametrization [8] as

$$\sin^2\theta_{12} = \frac{|U_{e2}|^2}{1-|U_{e3}|^2}\ , \qquad \sin^2\theta_{23} = \frac{|U_{\mu 3}|^2}{1-|U_{e3}|^2}\ , \qquad \sin\theta_{13} = |U_{e3}|\ . \qquad (75)$$



FIG. 1: The reactor mixing angle θ_{13} versus the ratio of third-to-second generation neutrino Yukawa couplings y_1^ν/y_2^ν (left plot) and the parameter $\kappa = y_R^\nu v_\chi/M$ (right plot). The (red) crosses and (blue) dots represent the results for the inverted and the normal mass hierarchy, respectively. The horizontal dotted (solid) lines in both plots indicate the upper and lower bounds on θ_{13} given in Eq. (1) at the 3σ level (best-fit value).

Leptonic CP violation at low energies can be detected through the neutrino oscillations which are sensitive to the Dirac CP-phase, but insensitive to the Majorana CP-phases in $U_{\rm PMNS}$ [22]: the Jarlskog invariant [20] is defined as

$$J_{CP} \equiv -{\rm Im}[U_{e1}^* U_{e3} U_{\tau 1} U_{\tau 3}^*] = \frac{1}{8}\sin 2\theta_{12}\sin 2\theta_{13}\sin 2\theta_{23}\cos\theta_{13}\sin\delta_{CP}\ , \tag{76}$$

where $U_{\alpha j}$ is an element of the PMNS matrix in Eq. (74), with $\alpha = e, \mu, \tau$ corresponding to the lepton flavors and $j = 1, 2, 3$ corresponding to the light neutrino mass eigenstates. And by manipulation of Eqs. (61) and (76) one can easily obtain the Dirac CP phase :

$$\delta_{CP} = -\arg\left(\frac{\frac{U_{e1}^* U_{e3} U_{\tau 1} U_{\tau 3}^*}{c_{12}c_{13}^2 c_{23} s_{13}} + c_{12}c_{23}s_{13}}{s_{12}s_{23}}\right)\ . \tag{77}$$

As expected, since the contributions of V_L^ℓ in Eq. (37) to the PMNS matrix are negligible, *i.e.*, its effects $\mathcal{O}(\lambda^3)$, we will consider $U_{\rm PMNS} \simeq U_\nu$ in numerical analysis.

IV. NUMERICAL ANALYSIS

Now we perform a numerical analysis for neutrinos using the linear algebra tools in Ref. [23].



FIG. 2: The reactor mixing angle θ_{13} versus the phase φ. The (red) crosses and (blue) dots represent the results for the inverted and normal mass hierarchy, respectively. The vertical dotted (solid) lines indicate the upper and lower bounds on θ_{13} given in Eq. (1) at the 3σ level (best-fit value).

The mass matrices $\widetilde{m}_D$ and $\widehat{M}_R$ in Eq. (62) contains seven parameters: $y^{\nu}_{3}, v_{\Phi}, M, y_1, y_2, \kappa, \varphi$. The first three ($y^{\nu}_{3}$, M and v_{Φ}) lead to the overall neutrino scale parameter m_0. The next four $(y_1, y_2, \kappa, \varphi)$ give rise to the deviations from TBM as well as the CP phases and corrections to the masse eigenvalues [see Eq. (64)]. Since we have a relation $v_{\chi}/\Lambda \sim \lambda^2$ in the charged fermion sector, for the cutoff scale $\Lambda = 10^{15}$ GeV we take $M = 10^{13}$ GeV and $v_{\Phi} = 172\sqrt{2}$ GeV, for simplicity, as inputs. Since the neutrino masses are sensitive to the combination $m_0 = v^2_{\Phi}|y^{\nu 2}_{3}|/(6M)$, other choices of M and v_{Φ} give identical results. Then the parameters $m_0, y_1, y_2, \kappa, \varphi$ can be determined from the experimental results of three mixing angles, $\theta_{12}, \theta_{13}, \theta_{23}$, and the two mass squared differences, $\Delta m^2_{\rm Sol}, \Delta m^2_{\rm Atm}$. In addition, the CP phases $\delta_{CP}, \varphi_{1,2}$ can be predicted after determining the model parameters. (Here, we will not discuss the Majorana CP phases $\varphi_{1,2}$.)

Using the formulae for the neutrino mixing angles and masses and our values of M, v_{Φ}, we obtain the following allowed regions of the unknown model parameters: for the normal mass hierarchy (NMH),

$$\begin{aligned} &0.21 \lesssim \kappa \lesssim 0.86\ , \qquad 1.04 \lesssim y_1 \lesssim 1.37\ , \qquad 1.04 \lesssim y_2 \lesssim 1.39\ , \\ &96^{\circ} \lesssim \varphi \lesssim 121^{\circ} \ \text{ and } \ 239^{\circ} \lesssim \varphi \lesssim 257^{\circ}\ , \qquad 1.1 \lesssim m_0 \times 10^{-2}[\text{eV}] \lesssim 3.6\ ; \end{aligned} \tag{78}$$



FIG. 3: The behaviors of θ_{23} and θ_{12} in terms of θ_{13}. The red crosses and the blue dots represent results for the inverted mass hierarchy and the normal mass hierarchy, respectively. The dotted vertical lines represent the experimental bounds of Eq. (1) at 3σ's.

for the inverted mass hierarchy (IMH),

$$\begin{aligned} &0.81 \lesssim \kappa \lesssim 2 \,, \qquad 0.91 \lesssim y_1 \lesssim 1.09 \,, \qquad 0.91 \lesssim y_2 \lesssim 1.08 \,, \\ &116^\circ \lesssim \varphi \lesssim 248^\circ \,, \qquad\qquad 1.6 \lesssim m_0 \times 10^{-2}[\mathrm{eV}] \lesssim 2.2 \,. \end{aligned} \tag{79}$$

Note that here we have used the 3σ experimental bounds on $\theta_{12}, \theta_{23}, \Delta m^2_{\rm Sol}, \Delta m^2_{\rm Atm}$ in Eq. (1), except for $\theta_{13} < 12^\circ$ for which we use the values in Eqs. (78,79). For these parameter regions, we investigate how mixing parameters do behave for the NMH and IMH. In Figs. 1-5, the data points represented by blue dots and red crosses indicate results for the NMH and IMH, respectively. The left-hand-side plot in Fig. 1 shows how the mixing angle θ_{13} depends on the ratio $y_1/y_2 = y^\nu_1/y^\nu_2$ of the third- and second-generation neutrino Yukawa couplings; the right-hand-side plot shows how θ_{13} depends on the parameter $\kappa = y^\nu_R v_\chi / M$. Fig. 2 shows the mixing angle θ_{13} as a function of the phase φ of $y^\nu_R v_\chi / M$. As can be seen in Figs. 1-2, only normal mass hierarchy is permitted within 3σ experimental bounds. And we see that the measured value of θ_{13} from the Daya Bay and RENO experiments can be achieved at 3σ's for $0.92 \lesssim y_1/y_2 < 1$, $1 < y_1/y_2 \lesssim 1.06$, $0.2 \lesssim \kappa \lesssim 0.7$, $95^\circ \lesssim \varphi \lesssim 115^\circ$ and $245^\circ \lesssim \varphi \lesssim 255^\circ$ for NMH. For IMH, in Figs. 1-2 the value of θ_{13} reaches at most 3°, which is excluded by the measurements of θ_{13}.

Fig. 3 shows how the values of θ_{13} depend on the mixing angles θ_{23} and θ_{12}. As can be seen in the left plot of Fig. 3, the behavior of θ_{23} in terms of the measured values of θ_{13} at



FIG. 4: The Jarlskog invariant J_{CP} versus the reactor angle θ_{13} (left plot), and the Dirac CP phase δ_{CP} versus θ_{23} (right plot). The (red) crosses and (blue) dots represent the results for the inverted and normal mass hierarchy, respectively. The vertical dotted (solid) lines indicate the upper and lower bounds on θ_{13} given in Eq. (1) at the 3σ level (best-fit value).

3σ's for the NMH is different than for the IMH. As already mentioned, the IMH is excluded by the measured values of θ_{13} in Fig. 3. For the NMH we see that the measured values of θ_{13} can be achieved for $49.5° \lesssim \theta_{23} \lesssim 54.8°$ and $35.8° \lesssim \theta_{23} \lesssim 40.5°$, with large deviations from maximality, which are favored at 1σ by the experimental bounds as can be seen in Eq. (1). Future precise measurements of θ_{23}, whether $\theta_{23} \to 45°$ or $|\theta_{23} - 45°| \to 5°$, will either exclude or favor our model. From the right plot of Fig. 3, we see that the predictions for θ_{13} do not strongly depend on θ_{12} in the allowed region.

To see how the parameters are correlated with low-energy CP violation observables measurable through neutrino oscillations, we consider the leptonic CP violation parameter defined by the Jarlskog invariant in Eq. (76) which can be expressed in terms of the elements of the matrix $h = \mathcal{M}_\nu \mathcal{M}_\nu^\dagger$ [22]:

$$J_{CP} = -\frac{\mathrm{Im}\{h_{12}h_{23}h_{31}\}}{\Delta m^2_{21}\Delta m^2_{31}\Delta m^2_{32}} \ . \tag{80}$$

The behavior of J_{CP} as a function of θ_{13} is plotted on the left plot of Fig. 4. We see that the value of J_{CP} lies in the ranges $0.006 \sim 0.03$ and $-0.03 \sim -0.004$ (NMH) for the measured value of θ_{13} at 3σ's. Also, in our model we have

$$\mathrm{Im}\{h_{12}h_{23}h_{31}\} = \frac{27 m_0^6}{4a^4b^3} y_1^2 y_2^2 (y_1^2 - y_2^2) \sin\psi_2 \{....\} \ , \tag{81}$$



FIG. 5: Plots of $|m_{ee}|$ as a function of θ_{13} and $m_{\rm lightest}$. The red crosses and the blue dots represent results for the inverted and the normal mass hierarchy, respectively. The vertical dashed lines show the experimental bounds of Eq. (1) at 3σ's.

in which $\{.....\}$ stands for a complicated lengthy function of y_1, y_2, a, b, ψ_1 and ψ_2. Clearly, Eq. (81) indicates that in the limit of $y_2 \to y_1$ or $\sin\psi_2 \to 0$ the leptonic CP violation J_{CP} goes to zero. When $y_2 \neq y_1$, i.e. for the IMH case, J_{CP} could go to zero as $\sin\psi_2$ of Eq. (81) [see, Eq. (52) and Fig. 2]. In the case of the NMH, J_{CP} has nonzero values for the measured range of θ_{13} while J_{CP} goes to zero for $\theta_{13} \to 0$, which corresponds to $y_2 \to y_1$. The right plot of Fig. 4 shows the behavior of the Dirac CP phase δ_{CP} [see Eq. (77)] as a function of θ_{23}, where the values of δ_{CP} lie in the ranges $0^\circ < \delta_{CP} \lesssim 60^\circ, 130^\circ \lesssim \delta_{CP} < 180^\circ, 180^\circ < \delta_{CP} \lesssim 240^\circ$ and $310^\circ \lesssim \delta_{CP} < 360^\circ$ for the NMH (for the IMH, δ_{CP} can vary over a wide range, but which is excluded by the measured values of θ_{13}). Interestingly, for the best-fit values of θ_{23} the values of δ_{CP} are predicted as the one around $10^\circ, 170^\circ, 190^\circ, 350^\circ$ for NMH. So, future precise measurements of θ_{23} will provide more information on δ_{CP}.

Moreover, we can straightforwardly obtain the effective neutrino mass $|m_{ee}|$ that characterizes the amplitude for neutrinoless double beta decay [25]:

$$|m_{ee}| \equiv \left| \sum_i (U_{\rm PMNS})^2_{ei} m_i \right| \,, \tag{82}$$

where $U_{\rm PMNS}$ is given in a good approximation as Eq. (69). The left and right plots in Fig. 5 show the behavior of the effective neutrino mass $|m_{ee}|$ in terms of θ_{13} and the lightest neutrino mass, respectively. In the left plot of Fig. 5, for the measured values of θ_{13} at 3σ's, the effective neutrino mass $|m_{ee}|$ can be in the range $0.04 \lesssim |m_{ee}|[\text{eV}] < 0.11$ for NMH. The right plot of Fig. 5 shows $|m_{ee}|$ as a function of m_{lightest}, where $m_{\text{lightest}} = m_1$ for the NMH and $m_{\text{lightest}} = m_3$ for the IMH. Our model predicts that the effective mass $|m_{ee}|$ is within the sensitivity about 10^{-2} eV of planned neutrinoless double-beta decay experiments [26].

V. CONCLUSION

Under $SU(2)_L \times U(1)_Y$ gauge symmetry, we have proposed a new model of leptons and quarks based on the discrete flavor symmetry T', the double covering of A_4. Here we impose that all Yukawa couplings be of order one, which implies that the hierarchies of charged fermion masses and the mildness of neutrino masses are responsible for six types of Higgs scalars. In addition to the gauge and flavor symmetries, in order to simplify our model and to remove the unwanted Yukawa terms appearing in the Lagrangian we have introduced a continuous global symmetry $U(1)_X$ which can not be gauged. After spontaneous $U(1)_X$ breaking, to avoid Goldstone bosons it has to be explicitly broken down to a subgroup Z_2. After spontaneous breaking of flavor symmetry, with the constraint of renormalizability in the Lagrangian, the leptons have $m_e = 0$ and the quarks have CKM mixing angles $\theta^q_{12} = 13^\circ, \theta^q_{23} = 0^\circ$ and $\theta^q_{13} = 0^\circ$. Thus, certain effective dimension-5 operators driven by the gauge-singlet and T'-triplet χ field are introduced as an equal footing, which induce $m_e \neq 0$ and lead the quark mixing matrix to the CKM one in form. On the other hand, the neutrino Lagrangian still keeps renormalizability. We have assumed that there is a cutoff scale Λ, above which there exists unknown physics.

We have shown numerical analysis in the lepton sector of our model, where only normal mass hierarchy is permitted within 3σ experimental bounds with the prediction of both large deviations from maximality in the atmospheric mixing angle θ_{23} and the measured values of reactor angle. So, future precise measurements of θ_{23}, whether $\theta_{23} \to 45^\circ$ or $|\theta_{23} - 45^\circ| \to 5^\circ$, would either exclude or favor our model. Together with it, our model has made predictions both for the Dirac CP phase $0^\circ < \delta_{CP} \lesssim 60^\circ, 130^\circ \lesssim \delta_{CP} < 180^\circ, 180^\circ < \delta_{CP} \lesssim 240^\circ$ and $310^\circ \lesssim \delta_{CP} < 360^\circ$, which is almost compatible with the global analysis in 1σ experimental bounds. Moreover, we have shown the effective mass $|m_{ee}|$ measurable in neutrinoless double beta decay to be in the range $0.04 \lesssim |m_{ee}|[eV] < 0.11$ for the normal hierarchy, which could be tested in near future neutrino experiments.

Appendix A: The Higgs potential



FIG. 6: Fifth dimension and locations of scalar and fermion fields.

In this Appendix, as an example, we present our Higgs potential and its minimization, as well as our prescription for effecting the stability of the vacuum alignment. We solve the vacuum alignment problem by extending the model into a spatial extra dimension y [24]. We assume that each field lives on a 4D brane either at $y = 0$ or at $y = L/2$ or at $y = L$, as shown in Fig. 6. The heavy neutrino masses and the neutrino Yukawa interactions arise from local operators at the branes $y = 0$ and $y = L/2$, while the charged fermion masses are realized by non-local effects involving both branes $y = 0$, $y = L/2$ and $y = L$. We impose that all the parameters appearing in the Lagrangian are assumed to be real. Once the scalars Φ, Ψ, χ, H, G acquire complex VEVs at the different branes, the CP symmetry can be spontaneously broken. A rigorous explanation of this possibility is beyond the scope of this paper.

The most general scalar potential ($d \leq 5$) for the Higgs fields Φ, H, Ψ, G, η and χ invariant under $SU(2)_L \times U(1)_Y \times T'$ and obeying the conditions in the previous paragraph, is then given by

$$V = V_{y=0} + V_{y=\frac{L}{2}} + V_{y=L}, \tag{A1}$$

where

$$\begin{aligned} V_{y=0} &= V(\Phi) + V(\Psi) + V(\Phi\Psi) \\ V_{y=\frac{L}{2}} &= V(\chi) + V(\eta) + V(\chi\eta) \\ V_{y=L} &= V(H) + V(G) + V(HG)\ , \end{aligned} \tag{A2}$$

and [7]

$$
\begin{aligned}
V(\Phi) &= \mu_\Phi^2(\Phi^\dagger\Phi)_{\mathbf{1}} + \lambda_1^\Phi(\Phi^\dagger\Phi)_{\mathbf{1}}(\Phi^\dagger\Phi)_{\mathbf{1}} + \lambda_2^\Phi(\Phi^\dagger\Phi)_{\mathbf{1}'}(\Phi^\dagger\Phi)_{\mathbf{1}''} \\
&+ \lambda_3^\Phi(\Phi^\dagger\Phi)_{\mathbf{3}_s}(\Phi^\dagger\Phi)_{\mathbf{3}_s} + \lambda_4^\Phi(\Phi^\dagger\Phi)_{\mathbf{3}_a}(\Phi^\dagger\Phi)_{\mathbf{3}_a} + \lambda_5^\Phi\{(\Phi^\dagger\Phi)_{\mathbf{3}_s}(\Phi^\dagger\Phi)_{\mathbf{3}_a} + h.c.\}\ , \qquad \text{(A3)} \\
V(\Psi) &= \mu_\Psi^2(\Psi^\dagger\Psi)_{\mathbf{1}} + \lambda_1^\Psi(\Psi^\dagger\Psi)_{\mathbf{1}}(\Psi^\dagger\Psi)_{\mathbf{1}} + \lambda_2^\Psi(\Psi^\dagger\Psi)_{\mathbf{1}'}(\Psi^\dagger\Psi)_{\mathbf{1}''} + \lambda_3^\Psi(\Psi^\dagger\Psi)_{\mathbf{3}_s}(\Psi^\dagger\Psi)_{\mathbf{3}_s} \\
&+ \lambda_4^\Psi(\Psi^\dagger\Psi)_{\mathbf{3}_a}(\Psi^\dagger\Psi)_{\mathbf{3}_a} + \lambda_5^\Psi\{(\Psi^\dagger\Psi)_{\mathbf{3}_s}(\Psi^\dagger\Psi)_{\mathbf{3}_a} + h.c.\}\ , \qquad \text{(A4)} \\
V(\Phi\Psi) &= \lambda_1^{\Phi\Psi}(\Phi^\dagger\Phi)_{\mathbf{1}}(\Psi^\dagger\Psi)_{\mathbf{1}} + \lambda_2^{\Phi\Psi}(\Phi^\dagger\Phi)_{\mathbf{1}'}(\Psi^\dagger\Psi)_{\mathbf{1}''} + \lambda_3^{\Phi\Psi}(\Phi^\dagger\Phi)_{\mathbf{3}_s}(\Psi^\dagger\Psi)_{\mathbf{3}_s} \\
&+ \lambda_4^{\Phi\Psi}(\Phi^\dagger\Phi)_{\mathbf{3}_a}(\Psi^\dagger\Psi)_{\mathbf{3}_a} + \lambda_5^{\Phi\Psi}\{(\Phi^\dagger\Phi)_{\mathbf{3}_s}(\Psi^\dagger\Psi)_{\mathbf{3}_a} + h.c.\} \\
&+ \lambda_6^{\Phi\Psi}\{(\Phi^\dagger\Phi)_{\mathbf{3}_a}(\Psi^\dagger\Psi)_{\mathbf{3}_s} + h.c.\} + \lambda_7^{\Phi\Psi}\left\{(\Phi^\dagger\Psi)_{\mathbf{1}}(\Phi^\dagger\Psi)_{\mathbf{1}} + h.c.\right\} \\
&+ \lambda_8^{\Phi\Psi}\left\{(\Phi^\dagger\Psi)_{\mathbf{1}'}(\Phi^\dagger\Psi)_{\mathbf{1}''} + h.c.\right\} + \lambda_9^{\Phi\Psi}\left\{(\Phi^\dagger\Psi)_{\mathbf{3}_s}(\Phi^\dagger\Psi)_{\mathbf{3}_s} + h.c.\right\} \\
&+ \lambda_{10}^{\Phi\Psi}\left\{(\Phi^\dagger\Psi)_{\mathbf{3}_a}(\Phi^\dagger\Psi)_{\mathbf{3}_a} + h.c.\right\} + \lambda_{11}^{\Phi\Psi}\left\{(\Phi^\dagger\Psi)_{\mathbf{3}_s}(\Phi^\dagger\Psi)_{\mathbf{3}_a} + h.c.\right\} \\
&+ \lambda_{12}^{\Phi\Psi}(\Phi^\dagger\Psi)_{\mathbf{1}}(\Psi^\dagger\Phi)_{\mathbf{1}} + \lambda_{13}^{\Phi\Psi}\left\{(\Phi^\dagger\Psi)_{\mathbf{1}'}(\Psi^\dagger\Phi)_{\mathbf{1}''} + h.c.\right\} + \lambda_{14}^{\Phi\Psi}(\Phi^\dagger\Psi)_{\mathbf{3}_s}(\Psi^\dagger\Phi)_{\mathbf{3}_s} \\
&+ \lambda_{15}^{\Phi\Psi}(\Phi^\dagger\Psi)_{\mathbf{3}_a}(\Psi^\dagger\Phi)_{\mathbf{3}_a} + \lambda_{16}^{\Phi\Psi}\{(\Phi^\dagger\Psi)_{\mathbf{3}_s}(\Psi^\dagger\Phi)_{\mathbf{3}_a} + h.c.\}\ , \qquad \text{(A5)}
\end{aligned}
$$

$$
\begin{aligned}
V(H) &= i\mu_H^2(H^\dagger H)_{\mathbf{1}} + \lambda_1^H(H^\dagger H)_{\mathbf{1}}(H^\dagger H)_{\mathbf{1}} + \lambda_2^H\{(H^\dagger H)_{\mathbf{3}}(H^\dagger H)_{\mathbf{3}} + h.c.\}\ , \qquad \text{(A6)} \\
V(G) &= i\mu_G^2(G^\dagger G)_{\mathbf{1}} + \lambda_1^G(G^\dagger G)_{\mathbf{1}}(G^\dagger G)_{\mathbf{1}} + \lambda_2^G\{(G^\dagger G)_{\mathbf{3}}(G^\dagger G)_{\mathbf{3}} + h.c.\}\ , \qquad \text{(A7)} \\
V(HG) &= \lambda_1^{HG}(H^\dagger H)_{\mathbf{1}}(G^\dagger G)_{\mathbf{1}} + \lambda_2^{HG}(H^\dagger G)_{\mathbf{1}}(G^\dagger H)_{\mathbf{1}} + \lambda_3^{HG}\{(G^\dagger H)_{\mathbf{1}}(G^\dagger H)_{\mathbf{1}} + h.c.\} \\
&+ \lambda_4^{HG}\{(H^\dagger H)_{\mathbf{3}}(G^\dagger G)_{\mathbf{3}} + h.c.\} + \lambda_5^{HG}\{(H^\dagger G)_{\mathbf{3}}(G^\dagger H)_{\mathbf{3}} + h.c.\} \\
&+ \lambda_6^{HG}\{(G^\dagger H)_{\mathbf{3}}(G^\dagger H)_{\mathbf{3}} + h.c.\}\ , \qquad \text{(A8)}
\end{aligned}
$$

$$
\begin{aligned}
V(\chi) &= \mu_\chi^2\left\{(\chi\chi)_{\mathbf{1}} + (\chi^*\chi^*)_{\mathbf{1}}\right\} + m_\chi^2(\chi\chi^*)_{\mathbf{1}} + \lambda_1^\chi\left\{(\chi\chi)_{\mathbf{1}}(\chi\chi)_{\mathbf{1}} + (\chi^*\chi^*)_{\mathbf{1}}(\chi^*\chi^*)_{\mathbf{1}}\right\} \\
&+ \lambda_2^\chi\left\{(\chi\chi)_{\mathbf{1}'}(\chi\chi)_{\mathbf{1}''} + (\chi^*\chi^*)_{\mathbf{1}'}(\chi^*\chi^*)_{\mathbf{1}''}\right\} \\
&+ \tilde\lambda_2^\chi\left\{(\chi^*\chi)_{\mathbf{1}'}(\chi\chi)_{\mathbf{1}''} + (\chi^*\chi)_{\mathbf{1}''}(\chi^*\chi^*)_{\mathbf{1}'}\right\} \\
&+ \lambda_3^\chi\left\{(\chi\chi)_{\mathbf{3}_s}(\chi\chi)_{\mathbf{3}_s} + (\chi^*\chi^*)_{\mathbf{3}_s}(\chi^*\chi^*)_{\mathbf{3}_s}\right\} + \tilde\lambda_3^\chi(\chi^*\chi)_{\mathbf{3}_s}\left\{(\chi\chi)_{\mathbf{3}_s} + (\chi^*\chi^*)_{\mathbf{3}_s}\right\} \\
&+ \lambda_4^\chi\left\{(\chi^*\chi)_{\mathbf{3}_a}(\chi\chi)_{\mathbf{3}_s} + (\chi\chi^*)_{\mathbf{3}_a}(\chi^*\chi^*)_{\mathbf{3}_s}\right\} \\
&+ \xi_1^\chi\left\{\chi(\chi\chi)_{\mathbf{3}_s} + \chi^*(\chi^*\chi^*)_{\mathbf{3}_s}\right\} + \tilde\xi_1^\chi\left\{\chi(\chi^*\chi^*)_{\mathbf{3}_s} + \chi^*(\chi\chi)_{\mathbf{3}_s}\right\}\ , \\
&+ \frac{\zeta_1^\chi}{\Lambda}\left\{(\chi\chi)_{\mathbf{1}}(\chi\chi\chi)_{\mathbf{1}} + (\chi^*\chi^*)_{\mathbf{1}}(\chi^*\chi^*\chi^*)_{\mathbf{1}}\right\} + \ldots + \frac{\zeta_{13}^\chi}{\Lambda}\{\ldots\}\ , \qquad \text{(A9)}
\end{aligned}
$$

---

[7] In $V(H)$ and $V(G)$ the terms $i\mu_H^2(H^\dagger H)_{\mathbf{1}}$ and $i\mu_G^2(G^\dagger G)_{\mathbf{1}}$ are expanded as $i\mu_H^2(H_1^\dagger H_2 - H_2^\dagger H_1)$ and $i\mu_G^2(G_1^\dagger G_2 - G_2^\dagger G_1)$, respectively.

$$
\begin{aligned}
V(\eta) &= \mu_{\eta}^{2}(\eta^{\dagger}\eta)+\lambda^{\eta}(\eta^{\dagger}\eta)^{2} \ , &\text{(A10)}\\
V(\chi\eta) &= \lambda_{1}^{\chi\eta}(\eta^{\dagger}\eta)\{(\chi\chi)_{\mathbf{1}}+(\chi^{*}\chi^{*})_{\mathbf{1}}\}\\
&+ \frac{\xi_{1}^{\chi\eta}}{\Lambda}(\eta^{\dagger}\eta)\{(\chi\chi\chi)_{\mathbf{1}}+(\chi^{*}\chi^{*}\chi^{*})_{\mathbf{1}}\}+\frac{\xi_{2}^{\chi\eta}}{\Lambda}(\eta^{\dagger}\eta)\{(\chi^{*}\chi\chi)_{\mathbf{1}}+(\chi\chi^{*}\chi^{*})_{\mathbf{1}}\} \ . &\text{(A11)}
\end{aligned}
$$

Here, $\mu_{\Phi}, \mu_{\Psi}, \mu_{H}, \mu_{G}, \mu_{\eta}, \mu_{\chi}$, m_{χ}, ξ_{1}^{χ}, $\tilde{\xi}_{1}^{\chi}$, $\zeta_{1\ldots13}^{\chi}$ and $\xi_{1,2}^{\chi\eta}$ have a mass dimension, whereas $\lambda_{1,\ldots,5}^{\Phi}$, $\lambda_{1,\ldots,5}^{\Psi}$, $\lambda_{1,2}^{H}$, $\lambda_{1,2}^{G}$, λ^{η}, $\lambda_{1,\ldots,4}^{\chi}$, $\tilde{\lambda}_{2,3}^{\chi}$, $\lambda_{1,\ldots,16}^{\Phi\Psi}$, $\lambda_{1,\ldots,4}^{HG}$ and $\lambda^{\chi\eta}$ are all dimensionless. And in $V(\chi)$ "$\cdots$" denotes dimension-5 operators composed of all possible combinations of χ fields.

1. Minimization of the neutral scalar potential

After the breaking of the flavor and electroweak symmetry, in the neutral Higgs sector, in order to find minimum configuration of the Higgs potential, we in general let

$$
\begin{aligned}
\langle\Phi_{i}\rangle &= \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}v_{\Phi_{j}}e^{i\gamma_{j}} \end{pmatrix} , \quad \langle\Psi_{j}\rangle = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}v_{\Psi_{j}}e^{i\zeta_{j}} \end{pmatrix} , \qquad \langle\eta\rangle = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}v_{\eta}e^{i\theta} \end{pmatrix} ,\\
\langle H_{k}\rangle &= \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}v_{H_{k}}e^{i\rho_{k}} \end{pmatrix} , \quad \langle G_{k}\rangle = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}v_{G_{k}}e^{i\sigma_{k}} \end{pmatrix} , \quad \langle\chi_{j}\rangle = v_{\chi_{j}}e^{i\varphi_{j}} \ , \qquad \text{(A12)}
\end{aligned}
$$

with $j=1-3, k=1,2$, where $v_{\Phi_{j}}, v_{\Psi_{j}}, v_{\eta}, v_{H_{k}}, v_{G_{k}}v_{\chi_{j}}$ are real and positive, and γ_{j}, ζ_{j}, $\rho_{k}, \sigma_{k}, \varphi_{j}$ are physically meaningful phases. Note that we can set $\theta=0$ without loss of generality because θ does not have physical meanings [see, Eqs. (A9)-(A11)]. First, at the brane $y=0$ the vacuum configuration for Φ and Ψ is obtained by vanishing of the derivative of V with respect to each component of the scalar fields Φ_{j} and Ψ_{j}. Then, we have six minimization equations for VEVs and six equations for phases. From those equations, we can get [8]

$$
\begin{aligned}
v_{\Phi}^{2} &\equiv v_{\Phi_{1}}^{2} = -\frac{18\mu_{\Phi}^{2}+v_{\Psi}^{2}W}{2(9\lambda_{1}^{\Phi}+4\lambda_{3}^{\Phi})} \neq 0 \ , \qquad \langle\Phi_{2}\rangle = \langle\Phi_{3}\rangle = 0 \ ,\\
v_{\Psi}^{2} &\equiv v_{\Psi_{1}}^{2} = -\frac{18\mu_{\Psi}^{2}+v_{\Phi}^{2}W}{2(9\lambda_{1}^{\Psi}+4\lambda_{3}^{\Psi})} \neq 0 \ , \qquad \langle\Psi_{2}\rangle = \langle\Psi_{3}\rangle = 0 \ , \qquad \text{(A13)}
\end{aligned}
$$

where $W=9\lambda_{1}^{\Phi\Psi}+4\lambda_{3}^{\Phi\Psi}+9\lambda_{12}^{\Phi\Psi}+4\lambda_{14}^{\Phi\Psi}+2(9\lambda_{7}^{\Phi\Psi}+4\lambda_{9}^{\Phi\Psi})\cos 2(\gamma-\zeta)$ with $\gamma\equiv\gamma_{1}$, $\zeta\equiv\zeta_{1}$, and v_{Φ} and v_{Ψ} are real. With the vacuum alignments of Φ,Ψ fields, Eq. (A13), minimal condition with respect to γ_{j},ζ_{j} are given as

$$
-\frac{\partial V}{\partial\gamma_{1}}\Big| = \frac{\partial V}{\partial\zeta_{1}}\Big| = \frac{v_{\Phi}^{2}v_{\Psi}^{2}}{9}\left(9\lambda_{7}^{\Phi\Psi}+4\lambda_{9}^{\Phi\Psi}\right)\sin 2(\gamma-\zeta) = 0 \ , \qquad \text{(A14)}
$$

---

[8] Of course, there are trivial solutions $v_{\Phi_{1}}=0, v_{\Psi_{1}}=0$. We have neglected them.

where $\gamma = \gamma_1,\ \zeta = \zeta_1$, and $\frac{\partial V}{\partial \gamma_{2,3}}\Big| = \frac{\partial V}{\partial \zeta_{2,3}}\Big| = 0$ is automatically satisfied.

Second, at the brane $y = L/2$ the vacuum configuration for χ and η is obtained by vanishing of the derivative of V with respect to each component of the scalar fields χ_j and η. For simplicity, we consider only the renormalizable terms in $V(\chi)$ and $V(\chi\eta)$. Then, we have seven minimization equations for four VEVs and three phases. From those equations, we can get [9]

$$v_{\chi_j}^2 \;=\; -\frac{m_\chi + 2(\mu_\chi^2 + v_\eta^2 \lambda^{\chi\eta})\cos 2\varphi}{12\{(\lambda_1^\chi + \lambda_2^\chi)\cos 4\varphi + \tilde{\lambda}_2^\chi \cos 2\varphi\}} = v_\chi^2 \neq 0 \ , \tag{A15}$$

where v_χ is real, and $\varphi_1 = \varphi_2 = \varphi_3 = \varphi$ is used. With the vacuum alignment of χ fields, Eq. (A15), minimal condition with respect to φ_i is given for $\varphi_1 = \varphi_2 = \varphi_3$ as

$$-\frac{1}{4}\frac{\partial V}{\partial \varphi_j}\Big| \;=\; v_\chi^2 \left\{ v_\eta^2 \lambda^{\chi\eta} + \mu_\chi^2 + 3v_\chi^2 \left(\tilde{\lambda}_2^\chi + 4(\lambda_1^\chi + \lambda_2^\chi)\cos 2\varphi_j \right) \right\} \sin 2\varphi_j = 0 \ , \tag{A16}$$

with $i = 1, 2, 3$. And, requiring vanishing of the derivative of V with respect to η,

$$\frac{1}{2}\frac{\partial V}{\partial \eta^0}\Big|^{<\chi>=v_\chi}_{<\eta^0>=v_\eta} \;=\; v_\eta \Big\{ v_\eta^2 \lambda^\eta + \frac{\mu_\eta^2}{2} + 3\lambda^{\Phi\chi} v_\chi^2 \cos 2\varphi \Big\} = 0 \ , \tag{A17}$$

and, we obtain the VEV of η for $\langle \chi \rangle = v_\chi e^{i\varphi}(1,1,1)$,

$$v_\eta^2 \;=\; \frac{-\mu_\eta^2 - 6v_\chi^2 \lambda^{\chi\eta}\cos 2\varphi}{2\lambda^\eta} \ . \tag{A18}$$

Finally, at the brane $y = L$ the vacuum configuration for H and G is obtained by vanishing of the derivative of V with respect to each component of the scalar fields H_i and G_i. Then, we have eight minimization equations for four VEVs and four phases. From those equations, we can get

$$\begin{aligned}
\frac{1}{2}\frac{\partial V}{\partial H_1^0}\Big|^{<G_j^0>=v_{G_j}}_{<H_j^0>=v_{H_j}} \;&=\; v_{H_1}\Big\{ 2v_{H_2}^2 \lambda_1^H (\cos 2\rho_{12} - 1) - \lambda_2^{HG} v_{G_2}^2 \Big\} + v_{H_2}\Big\{ \mu_H^2 \sin\rho_{12} \\
&-\; v_{G_1} v_{G_2} \left(2\lambda_1^{HG} \sin\sigma_{12}\sin\rho_{12} - \lambda_2^{HG}\cos(\sigma_{12} + \rho_{12}) \right) \Big\} = 0 \ , \\
\frac{1}{2}\frac{\partial V}{\partial H_2^0}\Big|^{<G_j^0>=v_{G_j}}_{<H_j^0>=v_{H_j}} \;&=\; v_{H_2}\Big\{ 2v_{H_1}^2 \lambda_1^H (\cos 2\rho_{12} - 1) - \lambda_2^{HG} v_{G_1}^2 \Big\} + v_{H_1}\Big\{ \mu_H^2 \sin\rho_{12} \\
&-\; v_{G_1} v_{G_2} \left(2\lambda_1^{HG} \sin\sigma_{12}\sin\rho_{12} - \lambda_2^{HG}\cos(\sigma_{12} + \rho_{12}) \right) \Big\} = 0 \ ,
\end{aligned} \tag{A19}$$

---

[9] There are trivial solutions $v_{\chi_j} = 0$. We have neglected them.

$$\begin{aligned}
\frac{1}{2}\frac{\partial V}{\partial G_1^0}\Big|^{<H_j^0>=v_{H_j}}_{<G_j^0>=v_{G_j}} &= v_{G_1}\Big\{2v_{G_2}^2\lambda_1^G(\cos 2\sigma_{12}-1)-\lambda_2^{HG}v_{H_2}^2\Big\}+v_{G_2}\Big\{\mu_G^2\sin\sigma_{12} \\
&- v_{H_1}v_{H_2}\left(2\lambda_1^{HG}\sin\sigma_{12}\sin\rho_{12}-\lambda_2^{HG}\cos(\sigma_{12}+\rho_{12})\right)\Big\}=0\ , \\
\frac{1}{2}\frac{\partial V}{\partial G_2^0}\Big|^{<H_j^0>=v_{H_j}}_{<G_j^0>=v_{G_j}} &= v_{G_2}\Big\{2v_{G_1}^2\lambda_1^G(\cos 2\sigma_{12}-1)-\lambda_2^{HG}v_{H_1}^2\Big\}+v_{G_1}\Big\{\mu_G^2\sin\sigma_{12} \\
&- v_{H_1}v_{H_2}\left(2\lambda_1^{HG}\sin\sigma_{12}\sin\rho_{12}-\lambda_2^{HG}\cos(\sigma_{12}+\rho_{12})\right)\Big\}=0\ . \quad \text{(A20)}
\end{aligned}$$

where $\rho_{12}\equiv\rho_1-\rho_2$ and $\sigma_{12}\equiv\sigma_1-\sigma_2$. And, we obtain the VEVs of H and G:

$$\begin{aligned}
v_{H_1} &= \frac{\mu_H^2\sin\rho_{12}-v_{G_1}v_{G_2}Y}{2v_{H_2}\lambda_1^H(1-\cos 2\rho_{12})+\lambda_2^{HG}v_{G_2}^2/v_{H_2}}\ , \\
v_{H_2} &= \frac{\mu_H^2\sin\rho_{12}-v_{G_1}v_{G_2}Y}{2v_{H_1}\lambda_1^H(1-\cos 2\rho_{12})+\lambda_2^{HG}v_{G_1}^2/v_{H_1}}\ , \\
v_{G_1} &= \frac{\mu_G^2\sin\sigma_{12}-v_{H_1}v_{H_2}Y}{2v_{G_2}\lambda_1^G(1-\cos 2\sigma_{12})+\lambda_2^{HG}v_{H_2}^2/v_{G_2}}\ , \\
v_{G_2} &= \frac{\mu_G^2\sin\sigma_{12}-v_{H_1}v_{H_2}Y}{2v_{G_1}\lambda_1^G(\cos 2\sigma_{12}-1)-\lambda_2^{HG}v_{H_1}^2/v_{G_1}}\ , \qquad \text{(A21)}
\end{aligned}$$

where $Y=2\lambda_1^{HG}\sin\sigma_{12}\sin\rho_{12}-\lambda_2^{HG}\cos(\sigma_{12}+\rho_{12})$. With the vacuum alignment of H and G fields, Eqs. (A19) and (A20), minimal condition with respect to ρ_j,σ_j is given as

$$\begin{aligned}
-\frac{1}{2}\frac{\partial V}{\partial\rho_1}\Big| = \frac{1}{2}\frac{\partial V}{\partial\rho_2}\Big| &= v_{H_1}v_{H_2}\Big\{-\mu_H^2\cos\rho_{12}+2v_{H_1}v_{H_2}\lambda_1^H\sin 2\rho_{12} \\
&+ v_{G_1}v_{G_2}\left(2\lambda_1^{HG}\sin\sigma_{12}\cos\rho_{12}+\lambda_2^{HG}\sin(\sigma_{12}+\rho_{12})\right)\Big\}=0\ , \quad \text{(A22)} \\
-\frac{1}{2}\frac{\partial V}{\partial\sigma_1}\Big| = \frac{1}{2}\frac{\partial V}{\partial\sigma_2}\Big| &= v_{G_1}v_{G_2}\Big\{-\mu_G^2\cos\sigma_{12}+2v_{G_1}v_{G_2}\lambda_1^G\sin 2\sigma_{12} \\
&+ v_{H_1}v_{H_2}\left(2\lambda_1^{HG}\sin\rho_{12}\cos\sigma_{12}+\lambda_2^{HG}\sin(\rho_{12}+\sigma_{12})\right)\Big\}=0\ . \quad \text{(A23)}
\end{aligned}$$

Appendix B

In Eq. (18) the components m^ℓ_{ij} are given by

$$\begin{aligned}
m^\ell_{11} &= Y_2^\mu\left(\frac{1-i}{2}\tilde{v}_{H_2}+i\tilde{v}_{H_1}\right)\ , \qquad m^\ell_{12}=Y_2^\mu\left(\frac{1-i}{2}\tilde{v}_{H_1}+\tilde{v}_{H_2}\right)\ , \\
m^\ell_{21} &= Y_1^\mu\left(\frac{1-i}{2}\tilde{v}_{H_2}+i\tilde{v}_{H_1}\right)\ , \qquad m^\ell_{22}=Y_1^\mu\left(\frac{1-i}{2}\tilde{v}_{H_1}+\tilde{v}_{H_2}\right)\ . \qquad \text{(B1)}
\end{aligned}$$

In Eq. (16) the components m^d_{ij} are given by

$$\begin{aligned}
m^d_{11} &= -Y^d_1\tilde{v}_{H_2} - Y^a_d\frac{1-i}{4}\tilde{v}_{H_2} + Y^s_d\left(\frac{2i}{3}\tilde{v}_{H_1} - \frac{1-i}{6}\tilde{v}_{H_2}\right) \\
m^d_{12} &= Y^d_1\tilde{v}_{H_1} + Y^a_d\left(\frac{1}{2}\tilde{v}_{H_2} - \frac{1-i}{4}\tilde{v}_{H_1}\right) - Y^s_d\left(\frac{1}{3}\tilde{v}_{H_2} + \frac{1-i}{6}\tilde{v}_{H_1}\right)\ ,\ m^d_{13} = \tilde{v}_\Psi\frac{2y^s_b}{3}\ , \\
m^d_{21} &= -Y^d_1\tilde{v}_{H_2} + Y^a_d\left(\frac{1-i}{4}\tilde{v}_{H_2} - \frac{i}{2}\tilde{v}_{H_1}\right) - Y^s_d\left(\frac{i}{3}\tilde{v}_{H_1} + \frac{1-i}{6}\tilde{v}_{H_2}\right) \\
m^d_{22} &= Y^d_1\tilde{v}_{H_1} + Y^a_d\frac{1-i}{4}\tilde{v}_{H_1} + Y^s_d\left(\frac{2}{3}\tilde{v}_{H_2} - \frac{1-i}{6}\tilde{v}_{H_1}\right)\ ,\quad m^d_{23} = -\tilde{v}_\Psi\left(\frac{y^a_b}{2} + \frac{y^s_b}{3}\right)\ , \\
m^d_{31} &= -Y^d_1\tilde{v}_{H_2} + Y^a_d\frac{i}{2}\tilde{v}_{H_2} + Y^s_d\left(\frac{1-i}{3}\tilde{v}_{H_2} - \frac{i}{3}\tilde{v}_{H_1}\right)\ , \\
m^d_{32} &= Y^d_1\tilde{v}_{H_1} - Y^a_d\frac{1}{2}\tilde{v}_{H_2} + Y^s_d\left(\frac{1-i}{3}\tilde{v}_{H_1} - \frac{1}{3}\tilde{v}_{H_2}\right)\ ,\ m^d_{33} = \tilde{v}_\Psi\left(\frac{y^a_b}{2} - \frac{y^s_b}{3}\right)\ .
\end{aligned} \tag{B2}$$

And, in Eq. (17) the components m^t_{ij} are given by

$$\begin{aligned}
m^t_{11} &= Y^a_u\left(\frac{1-i}{4}\tilde{v}_{G_2} - \frac{i}{2}\tilde{v}_{G_1}\right) - Y^s_u\left(\frac{1-i}{6}\tilde{v}_{G_2} + \frac{i}{3}\tilde{v}_{G_1}\right) \\
m^t_{12} &= Y^a_u\frac{(1-i)}{4}\tilde{v}_{G_1} + Y^s_u\left(\frac{2}{3}\tilde{v}_{G_2} - \frac{1-i}{6}\tilde{v}_{G_1}\right)\ ,\qquad m^t_{13} = \tilde{v}_\Phi\frac{2y^s_t}{3} \\
m^t_{21} &= Y^a_u\frac{i}{2}\tilde{v}_{G_1} + Y^s_u\left(-\frac{i}{3}\tilde{v}_{G_1} + \frac{1-i}{3}\tilde{v}_{G_2}\right) \\
m^t_{22} &= -Y^a_u\frac{1}{2}\tilde{v}_{G_2} + Y^s_u\left(\frac{1-i}{3}\tilde{v}_{G_1} - \frac{1}{3}\tilde{v}_{G_2}\right)\ ,\qquad m^t_{23} = -\tilde{v}_\Phi\left(\frac{y^a_t}{2} + \frac{y^s_t}{3}\right)\ , \\
m^t_{31} &= -Y^a_u\frac{1-i}{4}\tilde{v}_{G_2} + Y^s_u\left(\frac{2i}{3}\tilde{v}_{G_1} - \frac{1-i}{6}\tilde{v}_{G_2}\right)\ , \\
m^t_{32} &= Y^a_u\left(\frac{1}{2}\tilde{v}_{G_2} - \frac{1-i}{4}\tilde{v}_{G_1}\right) - Y^s_u\left(\frac{1-i}{6}\tilde{v}_{G_1} + \frac{1}{3}\tilde{v}_{G_2}\right)\ ,\quad m^t_{33} = \tilde{v}_\Phi\left(\frac{y^a_t}{2} - \frac{y^s_t}{3}\right)
\end{aligned} \tag{B3}$$

Acknowledgments

We thank prof. E. J. Chun for useful discussions.

[1] K. M. Case, R. Karplus and C. N. Yang, Phys. Rev. **101**, 874 (1956); P. H. Frampton and T. W. Kephart, Int. J. Mod. Phys. A **10**, 4689 (1995) [hep-ph/9409330]; A. Aranda, C. D. Carone and R. F. Lebed, Phys. Rev. D **62**, 016009 (2000) [hep-ph/0002044]; F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, Nucl. Phys. B **775**, 120 (2007) [Erratum-ibid. **836**, 127 (2010)] [hep-ph/0702194]; M. -C. Chen and K. T. Mahanthappa, Phys. Lett. B **652**, 34

(2007) [arXiv:0705.0714 [hep-ph]]; P. H. Frampton, T. W. Kephart and S. Matsuzaki, Phys. Rev. D **78**, 073004 (2008) [arXiv:0807.4713 [hep-ph]].

[2] E. Ma and G. Rajasekaran, Phys. Rev. D **64**, 113012 (2001) [arXiv:hep-ph/0106291]; K. S. Babu, E. Ma and J. W. F. Valle, Phys. Lett. B **552**, 207 (2003) [hep-ph/0206292]. G. Altarelli and F. Feruglio, Nucl. Phys. B **720**, 64 (2005) [arXiv:hep-ph/0504165] X. G. He, Y. Y. Keum and R. R. Volkas, JHEP **0604**, 039 (2006) [arXiv:hep-ph/0601001]. Y. H. Ahn and S. K. Kang, Phys. Rev. D **86**, 093003 (2012) [arXiv:1203.4185 [hep-ph]].

[3] M. -C. Chen, J. Huang, J. -M. O'Bryan, A. M. Wijangco and F. Yu, JHEP **1302**, 021 (2013) [arXiv:1210.6982 [hep-ph]]; M. Holthausen, M. Lindner and M. A. Schmidt, arXiv:1211.5143 [hep-ph]; S. F. King and C. Luhn, arXiv:1301.1340 [hep-ph].

[4] Y. H. Ahn, H. -Y. Cheng and S. Oh, Phys. Rev. D **83**, 076012 (2011) [arXiv:1102.0879 [hep-ph]]; Y. H. Ahn, C. S. Kim and S. Oh, arXiv:1103.0657 [hep-ph]; Y. H. Ahn, H. -Y. Cheng and S. Oh, Phys. Lett. B **715**, 203 (2012) [arXiv:1105.4460 [hep-ph]]; Y. H. Ahn, H. -Y. Cheng and S. Oh, Phys. Rev. D **84**, 113007 (2011) [arXiv:1107.4549 [hep-ph]]; Y. H. Ahn and H. Okada, Phys. Rev. D **85**, 073010 (2012) [arXiv:1201.4436 [hep-ph]]; S. Zhou, arXiv:1205.0761 [hep-ph]; S. Antusch, C. Gross, V. Maurer and C. Sluka, Nucl. Phys. B **866**, 255 (2013) [arXiv:1205.1051 [hep-ph]]; G. Altarelli, F. Feruglio and L. Merlo, arXiv:1205.5133 [hep-ph];

[5] Y. H. Ahn, S. Baek and P. Gondolo, Phys. Rev. D **86**, 053004 (2012) [arXiv:1207.1229 [hep-ph]].

[6] G. Altarelli and F. Feruglio, Nucl. Phys. B **741**, 215 (2006) [arXiv:hep-ph/0512103]; I. de Medeiros Varzielas, S. F. King and G. G. Ross, Phys. Lett. B **644**, 153 (2007) [arXiv:hep-ph/0512313]; G. Altarelli, F. Feruglio and Y. Lin, Nucl. Phys. B **775**, 31 (2007) [arXiv:hep-ph/0610165].

[7] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530**, 167 (2002); Z. Z. Xing, Phys. Lett. B 533, 85 (2002); P. F. Harrison and W. G. Scott, Phys. Lett. B **535**, 163 (2002); X. G. He and A. Zee, Phys. Lett. B 560, 87 (2003).

[8] J. Beringer et al. (Particle Data Group), Phys. Rev. D86, 010001 (2012).

[9] F. P. An *et al.* [DAYA-BAY Collaboration], Phys. Rev. Lett. **108**, 171803 (2012) [arXiv:1203.1669 [hep-ex]]; J. K. Ahn *et al.* [RENO Collaboration], Phys. Rev. Lett. **108**, 191802 (2012) [arXiv:1204.0626 [hep-ex]].

[10] K. Abe *et al.* [T2K Collaboration], Phys. Rev. Lett. **107**, 041801 (2011) [arXiv:1106.2822 [hep-

ex]]; see also: T. Nakaya [for the T2K Collaboration], talk at the Neutrino 2012 conference, http://neu2012.kek.jp/; P. Adamson *et al.* [MINOS Collaboration], Phys. Rev. Lett. **107**, 181802 (2011) [arXiv:1108.0015 [hep-ex]]; Y. Abe *et al.* [Double Chooz Collaboration], Phys. Rev. D **86**, 052008 (2012) [arXiv:1207.6632 [hep-ex]].

[11] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, JHEP **1212**, 123 (2012) [arXiv:1209.3023 [hep-ph]].

[12] J. Charles *et al.* [CKMfitter Group], Eur. Phys. J. C **41**, 1 (2005) [arXiv:hep-ph/0406184], and updated results from http://ckmfitter.in2p3.fr.

[13] M. Raidal, Phys. Rev. Lett. **93**, 161801 (2004) [arXiv:hep-ph/0404046]; H. Minakata and A. Y. Smirnov, Phys. Rev. D **70**, 073009 (2004) [arXiv:hep-ph/0405088]; F. Gonzalez Canales, A. Mondragon, AIP Conf. Proc. **857**, 287-292 (2006) [arXiv:hep-ph/0606175]; K. A. Hochmuth, W. Rodejohann, Phys. Rev. **D75**, 073001 (2007) [arXiv:hep-ph/0607103]; K. A. Hochmuth, S. T. Petcov, W. Rodejohann, Phys. Lett. **B654**, 177-188 (2007) [arXiv:0706.2975 [hep-ph]]; R. d. A. Toorop, F. Bazzocchi, L. Merlo, JHEP **1008**, 001 (2010) [arXiv:1003.4502 [hep-ph]]; J. Barranco, F. Gonzalez Canales, A. Mondragon, Phys. Rev. **D82**, 073010 (2010) [arXiv:1004.3781 [hep-ph]]; Y. Shimizu, R. Takahashi, [arXiv:1009.5504 [hep-ph]]; See also [4] : Y. H. Ahn, H. -Y. Cheng and S. Oh, Phys. Rev. D **83**, 076012 (2011).

[14] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986); G. F. Giudice *et al.*, Nucl. Phys. B **685**, 89 (2004) [arXiv:hep-ph/0310123]; W. Buchmuller, P. Di Bari and M. Plumacher, Annals Phys. **315**, 305 (2005) [arXiv:hep-ph/0401240]; A. Pilaftsis and T. E. J. Underwood, Phys. Rev. D **72**, 113001 (2005) [arXiv:hep-ph/0506107].

[15] A. Aranda, C. D. Carone and R. F. Lebed, Phys. Lett. B **474**, 170 (2000) [hep-ph/9910392].

[16] F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, Nucl. Phys. B **775**, 120 (2007) [Erratum-ibid. **836**, 127 (2010)] [hep-ph/0702194].

[17] J. C. Pati and A. Salam, Phys. Rev. Lett. **31**, 661 (1973); J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974) [Erratum-ibid. D **11**, 703 (1975)].

[18] R. Gatto, G. Sartori and M. Tonin, Phys. Lett. B **28**, 128 (1968); N. Cabibbo and L. Maiani, Phys. Lett. B **28**, 131 (1968); R. J. Oakes, Phys. Lett. B **29**, 683 (1969); H. Georgi and C. Jarlskog, Phys. Lett. B **86**, 297 (1979).

[19] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).

[20] C. Jarlskog, Phys. Rev. Lett. **55**, 1039 (1985); D. D. Wu, Phys. Rev. D **33**, 860 (1986).

[21] T. Fukuyama and H. Nishiura, arXiv:hep-ph/9702253; R. N. Mohapatra and S. Nussinov, Phys. Rev. **D 60**, 013002 (1999); E. Ma and M. Raidal, Phys. Rev. Lett. **87**, 011802 (2001); C. S. Lam, [arXiv:hep-ph/0104116]; T. Kitabayashi and M. Yasue, Phys.Rev. D **67** 015006 (2003); W. Grimus and L. Lavoura, arXiv:hep-ph/0305046; 0309050;W. Grimus and L. Lavoura, Phys. Lett. B **572**, 189 (2003); Y. Koide, Phys.Rev. D **69**, 093001 (2004); A. Ghosal, hep-ph/0304090; W. Grimus and L. Lavoura, J. Phys. G **30**, 73 (2004); R. N. Mohapatra and W. Rodejohann, Phys. Rev. D **72**, 053001 (2005) [hep-ph/0507312]; Y. H. Ahn, Sin Kyu Kang, C. S. Kim, Jake Lee, arXiv:hep-ph/0602160; Y. H. Ahn, S. K. Kang, C. S. Kim and J. Lee, Phys. Rev. D **75**, 013012 (2007).

[22] G. C. Branco, R. Gonzalez Felipe, F. R. Joaquim, I. Masina, M. N. Rebelo and C. A. Savoy, Phys. Rev. D **67**, 073025 (2003) [arXiv:hep-ph/0211001].

[23] S. Antusch, J. Kersten, M. Lindner, M. Ratz and M. A. Schmidt, JHEP **0503**, 024 (2005) [hep-ph/0501272].

[24] See [2] : G. Altarelli and F. Feruglio, Nucl. Phys. B **720**, 64 (2005).

[25] S. M. Bilenky, Phys. Part. Nucl. **41**, 690 (2010) [arXiv:1001.1946 [hep-ph]].

[26] C. Aalseth, H. Back, L. J. Dauwe, D. Dean, G. Drexlin, Y. Efremenko, H. Ejiri and S. Elliott *et al.*, hep-ph/0412300; I. Abt, M. F. Altmann, A. Bakalyarov, I. Barabanov, C. Bauer, E. Bellotti, S. T. Belyaev and L. B. Bezrukov *et al.*, hep-ex/0404039.